

**STUDIES ON TREMATODE PARASITES OF
CERTAIN FRESH WATER FISHES OF
BUNDELKHAND REGION**

THESIS

**Submitted for the Award of the Degree of
Doctor of Philosophy**

in

ZOOLOGY

**Under the Faculty of Science
of the**

Bundelkhand University, Jhansi


BY

Arvindra Kumar Yadav,

M. Sc.

**P. G. Department of Zoology
BIPIN BEHARI (P. G.) COLLEGE
JHANSI**

November 1995



T H E S I S

Submitted in fulfillment of the requirements

for the award of the degree of

DOCTOR OF PHILOSOPHY

in the subject

ZOOLOGY

Under the Faculty of Science

of

BUNDELKHAND UNIVERSITY

JHANSI

DR. U.K. DWIVEDI,
M. Sc., Ph. D.


P.G. Department of Zoology,
Bipin Behari(P.G.) College,
JHANSI.

Dated November, 1995.

CERTIFICATE

Certified that the thesis entitled, "Studies on Trematode Parasites of Certain Fresh-Water Fishes of Bundelkhand Region," submitted by Mr. Arvindra Kumar Yadav, M.Sc.(Zoology) for the degree of DOCTOR OF PHILOSOPHY (Ph.D.) in the subject ZOOLOGY under the Faculty of Science of the BUNDELKHAND UNIVERSITY, JHANSI embodies the original work done by him. He has worked under my guidance and supervision for more than twenty four months, commencing from the date of his registration.

It is further certified that the candidate has put in an attendance of over 200 days in this department from the date of his registration for the Ph.D. degree of the Bundelkhand University as required under relevant ordinances.



(U. K. DWIVEDI)
Supervisor

ACKNOWLEDGEMENTS

I wish to express my sincere gratitude and indebtedness to Dr. U.K. Dwivedi, M.Sc., Ph.D., Reader, P.G. Department of Zoology, Bipin Behari (P.G.) College, Jhansi, under whose valuable guidance and supervision the entire work embodied herein was planned and executed.

My grateful thanks are due to my Head of Department, Prof. J.P. Tewari, for his constant encouragement and interest in my work and for providing all facilities of laboratory and field work.

I owe a similar debt to my teacher Dr. S.C. Agarwal, Reader, P.G. Deptt. of Zoology, Bipin Behari (P.G.) College, Jhansi for extending all possible help to me in my work.

I wish to express my thanks to my friends and colleagues Mr. Arvind Kumar Yadav, Mr. Rahbar Sultan and Mr. Rajesh Kumar Dwivedi, for their kind co-operation and help in my work, specially during the collection of fishes from different places and in the laboratory work.

Lastly, I wish to express my gratefulness to Sri Rajbal Singh, Principal, G.N.K. Inter college, Jhansi for providing certain laboratory facilities in his college, and to Dr. O.P. Yadav, Reader, P.G. Deptt. of Zoology, Bipin Behari (P.G.) College, Jhansi for his valuable help in my daily laboratory work.

JHANSI,

Nov., 1995.

(Arvindra Kumar Yadav)

C O N T E N T S

	<u>Page No.</u>
Acknowledgement	
LIST OF CONTENTS	i - iv
 <u>PART - I</u>	
<u>GENERAL</u>	1 - 22
INTRODUCTION	1 - 5
HISTORICAL RESUME	6 - 12
MATERIAL AND METHOD	13 - 14
A SYSTEMATIC LIST OF HOSTS EXAMINED	15
HOST - PARASITE LIST	16 - 19
PARASITE - HOST LIST	20 - 22
 <u>PART - II</u>	
<u>MORPHOLOGY AND TAXONOMY OF CERTAIN TREMATODES</u>	23 - 70
Family : FELLODISTOMIDAE Nicoll, 1913	
Sub-family : FELLODISTOMINAE Nicoll, 1909	
Genus : <u>Pycnadena</u> Linton, 1911	
<u>Pycnadena bariliusi</u> Kumari, 1973	23 - 26
Family : ALLOCREADIIDAE Stossich, 1903	

Sub-family	:	ALLOCREADIINAE Looss, 1902	
Genus	:	<u>Allocreadium</u> Looss, 1900	
		<u>Allocreadium handiai</u>	
		Pande, 1937	27 - 30
Sub-family	:	ORIENTOCREADIINAE Yamaguti, 1958	
Genus	:	<u>Orientocreadium</u>	
		Tubangui, 1931	
		<u>Orientocreadium batrachoides</u>	
		Tubangui, 1931	31 - 33
		<u>Orientocreadium indicum</u>	
		Pande, 1934	34 - 37
		<u>Orientocreadium pseudobagri</u>	
		Yamaguti, 1934	38 - 40
		<u>Orientocreadium keni</u> n.sp.	41 - 43
Family	:	GORGODERIDAE Looss, 1901	
Sub-family	:	PHYLLODISTOMINAE Yamaguti, 1958	
Genus	:	<u>Phyllodistomum</u> Braun, 1899	
		<u>Phyllodistomum tripathi</u>	44 - 47
		Motwani and Srivastava, 1961	
Family	:	ISOPARORCHIIDAE Poche, 1926	
Genus	:	<u>Isoparorchis</u>	
		Southwell, 1913	
		Metacercaria of	
		<u>Isoparorchis hypselobagri</u>	48 - 49
		(Billet, 1898) Odhner, 1911	
Family	:	HEMIURIDAE Luhe, 1901	
Sub-family	:	HALIPEGINAE Ejsmont, 1931	

Genus	:	<u>Genarchopsis</u> Ozaki, 1925	
		<u>Genarchopsis piscicola</u> Srivastava, 1933	50 - 53
		<u>Genarchopsis goppo</u> Srivastava, 1933	54 - 56
		<u>Genarchopsis singularis</u> Srivastava, 1933	57 - 60
Family	:	CLINOSTOMIDAE Luhe, 1901	
Genus	:	<u>Clinostomum</u> Leidy, 1856	
		Metacercaria of <u>Clinostomum Complanatum</u> (Rudolphi, 1819) Braun, 1899	61
Sub-family	:	EUCLINOSTOMINAE Yamaguti, 1958	
Genus	:	<u>Euclinostomum</u> Travassos, 1928	
		Metacercaria of <u>Euclinostomum heterostomum</u> (Rudolphi, 1809) Travassos, 1928.	62 - 66
Family	:	HETEROPHYIDAE Odhner, 1914	
Sub-family	:	HAPLORCHIINAE Looss, 1899	
Genus	:	<u>Haplorchoides</u> Chen, 1949	
		<u>Haplorchoides seenghali</u> Dayal and Gupta, 1954	67 - 70

PART - III

<u>HOST-PARASITE RELATIONSHIP AND SEASONAL INCIDENCE</u>	71 - 108
- Host-Parasite relationship	71 - 73
- Index of total helminth infection (Trematodes)	74 - 75
- Host-wise analysis of parasites	76 - 83
- Overall incidence of Trematodes	84 - 89
- Level and intensity of parasitization	90 - 100
- Seasonal incidence of parasites	101 - 108
 EXPLANATION OF LIST/TABLES	 109 - 110
 EXPLANATION OF PLATES AND FIGURES (PLATES AND FIGURES 1 TO 19)	 111 - 113
 SUMMARY	 114 - 121
 REFERENCES	 122 - 173

- - - - -

P A R T - I

INTRODUCTION

Fishes occupy an important place in human diet, economy and ecosystem. Like other vertebrates, fishes too harbour parasitic population of diverse group of helminth parasites, which have detrimental effect on the fishes in many ways like stunted growth, postponement of sexual maturity, damage to various organs thus affecting the yield of fish production. The study of fish pathology is of much importance both from the point of view of fishery management and also to check the spread of human and animal diseases for which fishes act as carriers. The success of implementation of various fishery development programmes depends to a certain extent on the intensification of fish parasitological research, as the improvement of fish production can be achieved from healthy fish stock only.

During the last five decades, lot of work has been done on the morphology and taxonomy of parasites of fishes. These researches have contributed a lot to our knowledge of many genera and species, both known and new. Still our knowledge on fish parasites and disease is very meagre and we have no knowledge of the amount of damage caused and loss incurred in total fish production and fish products due to parasites and parasitic diseases.

In Uttar Pradesh and Madhya Pradesh too, ambitious projects are being launched to develop inland fisheries. But little work has been done on the parasites of fishes particularly from Bundelkhand region. Keeping this in view, the present study was undertaken and attempt has been made to fill a part of the wide lacuna left in this field.

Bundelkhand region is considered to be a premier region of the country, because it has a number of small and large water bodies, lakes, dams and two rivers which provide better fisheries and aquaculture prospects. All these water bodies are rich in fish fauna.

The Bundelkhand region forms South-east boundry of Uttar Pradesh, extending from 24.21' to 26.42' North latitude and 78.14' to 81.38' East longitude. It is comprised of five districts, namely - Jhansi, Lalitpur, Jalaun, Hamirpur and Banda. The region is surrounded in northern side by the districts of Etawah, Kanpur, Fatehpur and Allahabad of Uttar Pradesh; in Western side by the districts of Guna, Shivpuri and Datia of Madhya Pradesh, and in Southern side by the districts of Sagar, Chattarpur Panna of Madhya Pradesh.

The river Betwa and river Ken are two of the major rivers of the region. In the Bundlkhand region a number of large and small dams are also present which are a good source of inland fisheries. The main dams are - Jamni dam, Sagnam dam, Rohni dam, Rjghat dam, Matatila dam, Dhukwan dam, Shahzad dam, Saprar dam, Pahuji dam, Khaprar dam, Dongri dam, Maudaha dam, Arjun dam, Kabrai dam, Barwa dam, Gunta dam, Pathra dam, Parichha dam, Pahari dam, Keolari dam, Chandrawal dam. A number of rivulets, large number of ponds, pools and irrigation canals and drains constitute considerably rich fresh-water resources for inland fisheries. The daily fish-output from this region is 30 to 40 metric tons in off season and 50 to 80 metric tons during the season. The whole collection is exported to different important areas of the country after satisfying the regional requirements. All popular groups of fishes such as major carps, cat fishes, live fishes, feather backs, sheet fishes, eels etc. form the bulk of total production of this region.

The present work pertains to a group of air-breathing fishes, commonly available in fresh water bodies of the region namely Clarias batrachus (Linn.), Heteropneustes fossilis (Bl.) and species of Channa. These include Channa punctatus (Bl.), Channa striatus (Bl.) and Channa marulius (Ham.). Out of the five species of Channa available in Madhya Pradesh, Channa gachua (Ham.), Channa marulius (Ham.)

Channa punctatus (B1.), Channa striatus (B1.) and Channa leucopunctatus (B1.), only the above mentioned three species are available in the water dams. These fishes are predatory in nature. Since these fishes remain alive for a long period out of water, so they are usually marketed alive and also known as live fishes. These fishes are easy to handle in laboratory. Though these fishes are mostly non-commercial, yet they have their own economic value and great demand due to their high protein, high iron and low fat contents compared to that in carps. Clarias batrachus (Linn.) and H. fossilis (B1.) are considered to be highly nutritious and esteemed as food where as species of Channa are eaten mostly by poor classes.

Thus efforts have been made to concentrate the work on these host fishes and to obtain maximum number of parasites from them throughout the period of study.

In order to make the faunistic studies more elaborate and objective, host parasite relationship has also been studied and analysed statistically, in order to study index of total helminth infection; host-wise analysis; overall incidence; level and intensity of parasitization; seasonal incidence; and intensity of infection in trematodes. Such calculations can help considerably in the correct understanding of the nature and extent of their pathogenic role.

The present findings definitely contribute to the general survey of the parasitic fauna of this region. It may help the pisciculturists in understanding better the effect of parasitization on the fishes, thus helping to increase the production of fish as food to human beings.

- - - - -

HISTORICAL RESUME

HISTORICAL RESUME

Helminth parasites have been known to mankind since Vedic and Post-Vedic period around 800 B.C., as in Atharvaveda these worms are referred to as Krimis. Great medical treatises of Charaka and Susruta (between 200 B.C. and 200 A.D.) even mentioned detailed treatment of parasites. A detailed account of knowledge of these worms in ancient India is given by Bhaduri, Tiwari and Biswas (1972).

Our present day knowledge of helminth parasites dates back to 1379, when Jean - de - Brie (1379) who for the first time discovered fluke Fasciola. The first references to trematodes, probably Fasciola jacksoni and Pseudodyscus hawkesi, from Indian region in modern times is made by Gilchrist, who has worked on them in the years 1841-1846. Later on, Cobbold (1869-1882) wrote a series of papers describing parasites of elephant, cattle and Gangeti dolphin and thus making the beginning of the scientific study of trematode fauna of India.

Bhalerao (1926) was the first Indian helminthologist to give a boost to Indian helminthology in general and study of trematodes in particular. He was followed by a band of dedicated workers like Mehra, Verma, Moghe, Thapar, Lal, Srivastava (H.D.), Chauhan, Pande and many others, who contributed much to our present day knowledge of the

trematodes from Indian region. Bhalerao (1939) reviewed the progress of the knowledge of trematodes in India till that time. In 1956, Thapar and Chauhanⁱⁿ 1963 also discussed the progress of helminthology in India with special reference to trematodes. Some of the more important contributions in this field include -

Bhalerao (1926, 36); Verma (1927, 36); Chauhan (1940, 49, 54, 55); Srivastava, H.D. (1938, 48); Mehra, H.R. (1935, 38, 62, 66); Pande (1937); Patwardhan (1935); Khan (1935); Mehra, R.K. (1941); Kaw (1950); Dayal (1948, 49, 50); Baugh (1949, 50); Gupta (1950, 51, 55, 56); Jaiswal (1957), Jain (1967) and Pandey, K.C. (1970).

Helminth parasites of Indian fishes have not received the attention they deserve, except for the systematics of certain parasites. A perusal of literature shows that following Indian workers have described the trematode parasites of fishes -

Billet (1899) described Isoparorchis hypselobagri from Wallagoattu and also immature forms of the same from Barbus tor, Channa striatus, Notopterus notopterus and Mastacembelus armatus from India.

Southwell (1913) described Isoparorchis trisimilitubis which was later renamed as I. hypselobagri from the air bladder of wallago attu. Verma (1927) reported Opisthorchis

pedicellata from the gall bladder of Rita rita. He (1935) also recorded gastorostones from siluroid fishes.

A large number of workers have made substantial contribution on the taxonomy of trematode parasites. These include :-

Southwell and Prashad (1918) described Clinostomum piscidum from Nandus nandus.

Verma (1927) described Opisthorchis pedicellata from Bagarius varrellii and Rita rita from Allahabad.

Thapar (1930) described Gomtia piscicola from Bagarius varrellii from Lucknow.

Srivastava, H.D. (1933) described Progonus piscicola and Progonus ovocaudata from Ophiocephalus punctatus; Ophiocorchis lobata and Ophiocorchis singularis from Ophiocephalus striatus from Allahabad.

Pande (1934) described Orientocreadium indicum from Heteropneustes fossilis and Rita buehanani from Allahabad. In (1937) he described Allocreadium handiai from Ophiocephalus punctatus. In (1938) he described Allocreadium kosia from Barbus Chilinoidea; Allocreadium schizothoracis from Schizothorax micropogon; Allocreadium mahaseri from Barbus tor from Allahabad.

Dayal (1949) described Phyllodistomus vachius from Eutropiichthys vacha from Lucknow and Allahabad.

Bhalerao (1941) described Clinostomum indicum from Notopterus notopterus from Allahabad. In (1942) he described Clinostomum dasi from Saccobranchus fossilis and Clinostomum prashadi from an unidentified fish from Hyderabad.

Mehra (1941) described Opisthorchis pedicellata minutus from Mystus seenghala and Wallago attu from Allahabad.

Gupta (1950) described Allocreadium thapari from Rita rita from Hardoi. In (1951) he described Phyllodistomum singhiai from Mastacembelus armatus from Lucknow and Saharanpur. In (1953) he described Haplorchoides seenghali from Macrones seenghala; Phyllodistomum vittatusi from Macrones vittatus; Haplorchoides ritai, Haplorchoides brahamputraensis from Rita rita from Assam; Haplorchoides gontioensis from Silundia gangetica from Lucknow. In (1956) he described Allocreadium kamali from Chela bacaila, Allocreadium mehrai from Rhychobdella aculeata from Lucknow. In (1963) he described Allocreadium makundai from Barbus sarana from Banaras.

Gupta and Verma (1976 Publ. 1977) described Allocreadium mrigali, Allocreadium baranai, Allocreadium saranai from Cirrhina mrigala, Barilius barana and Barbus sarana respectively.

Kaw (1950) described Allocreadium nemacheilus from Nemacheilus kashmirensis; Clinostomum schizothoroxi from Oreinus sinatus, Schizothorax micropogon; Phyllodistomum loossi from Schizothorax socinus from Kashmir.

Jaiswal (1957) described Phyllodistomum (Catroptoides) indianum from Heteropneustes fossilis and Phyllodistomum Parorchium from Glossogobius (Gobius) giuris; Euclinostomum chanai from Ophiocephalus punctatus; Clinostomum macrosomium from Mastacembelus armatus from Hyderabad.

Saksena (1958) described Orientocreadium raipurensis, Orientocreadium dayali from Clarias batrachus; Allocreadium spindala from Mastacembelus armatus from Raipur. In (1960) he described Orientocreadium umadasi from Clarias batrachus from Raipur.

Srivastava, P.S. (1960) described Allocreadium ophiocephali from Ophiocephalus punctatus from Raipur.

Motwani and Srivastava (1961) described Phyllodistomum chauhani from Mystus tor and Mystus seenghala; Phyllodistomum tripathi from Bagarius bagarius from India.

Rai (1962) described Allocreadium dollfusi, Allocreadium singhi, Allocreadium hirnai from Barbus tor from the River Hiran, near Katangi and Sehora (M.P.).

Srivastava, C.B. (1962) described Pycnadena komiyai from Oxygaster gora from India.

Agrawal (1964) described Allocreadium heteropneustus from Heteropneustes fossilis; Haplorchoides macroni from Macrones seenghala from Lucknow. In (1966) she described Genarchopsis punctati from Ophiocephalus punctatus from Lucknow.

Kakaji (1969) described Genarchopsis cameroni from Mystus seenghala; Allocreadium catlai from Catla catla; Genarchopsis cuchiai from Amphipnous cuchia from Lucknow. In the same year, she described Allocreadium guptai and Allocreadium fasciatusi from Rita rita and Trichoquaster fasciatus, respectively from Varanasi.

Fotedar (1969) described Phyllodistomum megacotyle from Garra mullya from Kashmir.

Pande, B.P. and Shukla, R.P. (1976) described Haplorchoides pearsoni and Haplorchoides mehrai from Channa punctatus and Mystus vittatus, respectively from Lucknow.

Gupta, V. and Puri, M. (1979, Publ. 1980) described Allocreadium calbassii, Allocreadium manteri from Labeo calbasu, Anabas testudineus, respectively from Lucknow.

Systematics of fish cestodes of India region have been studied by Hornell (1912); Southwell (1913, 1915, 1925, 1930); Moghe (1925); Woodland (1924); Subramaniam (1939, 1940) and many other workers. Johri (1959) described new cestodes from fishes. Rao (1960) studied Mehlis gland complex by histochemical methods and also described a new species of *ptychobothrium* from a flying fish. Agarwal (1965-1980) studied fish cestodes from Raipur. Rajyalaxmi (1981-1988) studied Cestodes of Waltair Coast fishes.

In Madhya Pradesh also, lot of work was done at Jabalpur, Raipur, Rewa, Ujjain and Gwalior by Singh, Agarwal, Dwivedi, Rai, Saxena, Jain, Khoche, Johri, Dandotia and others. However little work has been done on the helminth parasites of fishes of Bundelkhand region.

From the foregoing account, it is evident that considerable progress is being made in the knowledge of helminth fauna and its taxonomic study of this country, but very little work has been done to ascertain the incidence of parasites and estimation of helminthic infection. The important contributions in this field have been made by Srivastava, C.B. and Mukherjee, G.D. 1986; Siddiqui A.H.; Nizami, W.A. (1988) Devraj, M & Ranganathan, V (1991); Bahaduria, S. (1982), Dandotia, M.R. (1984, 1992, 1994) and others. Still a wide lacuna is left in this field considering the richness of fauna of parasite helminthes.

MATERIAL AND METHOD

MATERIAL AND METHOD

The host fishes were collected from several dams, ponds and other local fresh water bodies in and around Jhansi and in the Bundelkhand region. Fishes were also purchased from local fish markets of Sipri and Raiganj Bazar.

These host fishes were kept alive in aquaria in the laboratory and then freshly killed, dissected and examined at convenience. A thorough search was done to determine the where-abouts of parasites. Various organs particularly body cavity, stomach, duodenum, intestine, rectum, gall bladder and Kidneys were carefully examined in petridish under low power binocular.

Soon after collection, the trematodes were thoroughly washed and kept in saline water. They were studied alive and observations were made regarding the colour and movements of body, spines on the body, oral and ventral suckers, cirrus and metraterm, excretory bladder and its branches.

For fixation 5-10% formaline was used. For whole mounts, preservation in formaline for longer periods gave good results. For preparing whole mounts, precaution was taken to avoid over or under pressure.

For preparing whole mount of trematodes, after fixation and thorough washing in water, worms were dehydrated and stained in Borox Carmine, then cleared in xylene and finally mounted in DPX. The drawings of the whole mounts were made with the help of camera lucida at a suitable magnification.

To record the seasonal incidence, the fishes were examined at regular intervals from July 1993 to June 1995. In all, 75 fishes per month with an average of 15 fishes per month of each species were examined. A total of 1800 fishes were collected and studied during two years period. The data collected was studied and statistically analysed. Thus index of total helminth infection (Trematodes), host-wise analysis, overall incidence, level and intensity of parasitization, seasonal incidence.

- - - - -

A SYSTEMATIC LIST OF THE HOSTS EXAMINED

A SYSTEMATIC LIST OF THE HOSTS EXAMINED

Class	Teleostomi
Subclass	Actinopterygii
Order	Cypriniformes
Division	Siluri
Sub-order	Siluroidei
Family	Clariidae
Example	<u>Clarias</u> <u>batrachus</u> (Linn.)
Family	Saccobranchidae
Example	<u>Heteropneustes</u> <u>fossilis</u> (B1.)
Order	Ophiocephaliformes
Family	Ophiocephalidae
Example	<u>Channa</u> <u>punctatus</u> (B1.)
Example	<u>Channa</u> <u>striatus</u> (B1.)
Example	<u>Channa</u> <u>marulius</u> (Ham.)

HOST - PARASITE LIST

HOST - PARASITE LIST

Selected fresh water fishes have been examined for the present study of helminth parasites. These fishes are - Clarias batrachus (Linn.), Heteropneustes fossilis (B1.) and sp. of Channa - namely : Channa punctatus (B1.) Channa striatus (B1.) and Channa marulius (Ham.) which ^{and in Bundelkhand region} are available locally in plenty. These live fishes were examined thoroughly for helminth infection and parasites were collected.

The following host - parasite list includes only the forms collected by the author and described in the present thesis. This includes species of digenetic trematodes only. Majority of the parasites form first host and locality record from this region and a new species recovered from Clarias batrachus and Heteropneustes fossilis.

Host - Parasite list

<u>Name of the host</u>	<u>Name of the parasite</u>
<u>Channa punctatus</u> (B1.)	<u>Genarchopsis singularis</u> Srivastava, 1933.
	<u>Genarchopsis goppo</u> Srivastava, 1933
	<u>Genarchopsis piscicola</u> Srivastava, 1933.

Orientocreadium indicum

Pande, 1934.

Allocreadium handiai

Pande, 1937.

Metacercaria

Euclinostomum heterostomum

(Rudolphi, 1809) Travassos, 1928.

Isoparorchis hypselobagri

(Billet, 1898) Odhner, 1911.

Clinostomum complanatum

(Rudolphi, 1819) Braun, 1899.

Channa striatus (Bl.)Genarchopsis singularis

Srivastava, 1933.

Orientocreadium indicum

Pande, 1934.

Allocreadium handiai

Pande, 1937.

Metacercaria

Isoparorchis hypselobagri

(Billet, 1898) Odhner, 1911.

Channa marulius (Ham.)Orientocreadium indicum

Pande, 1934

Metacercaria

Isoparorchis hypselobagri

(Billet, 1898) Odhner, 1911.

Heteropneustes fossilis

(B1.)

Orientocreadium indicum

Pande, 1934.

Orientocreadium pseudobagri

Yamaguti, 1934.

Orientocreadium keni n. sp.Allocreadium handiai

Pande, 1937

Haplorchoides seenghali

Dayal and Gupta, 1954.

Phyllodistomum tripathi

Motwani and Srivastava, 1961.

Clarias batrachus (Linn.)Orientocreadium indicum

Pande, 1934

Orientocreadium Pseudobagri

Yamaguti, 1934.

Orientocreadium keni n. sp.Orientocreadium batrachoides

Tubangui, 1931.

Allocreadium handiai

Pande, 1937.

Pycnadena bariliusi

V. Kumari, 1973.

PARASITE - HOST LIST

PARASITE - HOST LIST

<u>Parasite</u>	<u>Host</u>	<u>Location</u>
<u>Pycnadena bariliusi</u> V. Kumari, 1973.	<u>Clarias batrachus</u> (Linn.)	Stomach
<u>Allocreadium handiai</u> Pande, 1937.	<u>Clarias batrachus</u> (Linn.) <u>Heteropneustes fossilis</u> (Bl.) <u>Channa punctatus</u> (Bl.) <u>Channa striatus</u> (Bl.)	Intestine Intestine Intestine Intestine
<u>Orientocreadium batrachoides</u> Tubangui, 1931.	<u>Clarias batrachus</u> (Linn.)	Intestine
<u>Orientocreadium indicum</u> Pande, 1934	<u>Clarias batrachus</u> (Linn.) <u>Heteropneustes fossilis</u> (Bl.) <u>Channa punctatus</u> (Bl.) <u>Channa striatus</u> (Bl.) <u>Channa marulius</u> (Ham.)	Intestine Intestine Intestine Intestine Intestine
<u>Orientocreadium pseudobagri</u> Yamaguti, 1934.	<u>Clarias batrachus</u> (Linn.)	Intestine

<u>Parasite</u>	<u>Host</u>	<u>Location</u>
<u>Orientocreadium keni</u> n.sp.	<u>Clarias batrachus</u> (Linn.)	Intestine
	<u>Heteropneustes fossilis</u> (B1.)	Intestine
<u>Phyllodistomum tripathi</u> Motwani and Srivastava, 1961.	<u>Heteropneustes fossilis</u> (B1.)	Intestine
<u>Isoparorchis hypselobagri</u> (Billet, 1898) Odhner, 1911	<u>Channa punctatus</u> (B1.)	Body cavity
	<u>Channa striatus</u> (B1.)	Body cavity
	<u>Channa marulius</u> (B1.)	Body cavity
<u>Genarchopsis piscicola</u> Srivastava, 1933.	<u>Channa punctatus</u> (B1.)	Intestine
<u>Genarchopsis goppo</u> Srivastava, 1933.	<u>Channa punctatus</u> (B1.)	Intestine
<u>Genarchopsis singularis</u> Srivastava, 1933	<u>Channa punctatus</u> (B1.)	Intestine
	<u>Channa striatus</u> (B1.)	Intestine

<u>Parasite</u>	<u>Host</u>	<u>Location</u>
<u>Euclinostomum heterostomum</u> (Rudolphi, 1809) Travassos, 1928.	<u>Channa punctatus</u> (Bl.)	Body cavity
<u>Clinostomum complanatum</u> (Rudolphi, 1819) Braun, 1899.	<u>Channa punctatus</u> (Bl.)	Body cavity
<u>Haplorchoides seenghali</u> Dayal and Gupta, 1954.	<u>Heteropneustes fossilis</u> (Bl.)	Intestine

P A R T - I I

Ovary pretesticular and intertesticular, submedian, diagonally oval, measures 0.17×0.010 . Receptaculum seminis large. Uterus very extensive, ascending and descending coils extend from pharynx to posterior end of the body. Eggs numerous, oval, yellow, small. Vitelline follicles lateral, extending from oral sucker to posterior end of the body, confluent in posttesticular region.

Discussion

Linton (1911) proposed the name pyncadena to replace the generic name Didymorchis erected by him in 1910, since it was preoccupied. Srivastava, C.B. (1962) described Pyncadena komiyai and placed it under the family Allocreadiidae following Yamaguti (1958), Manter (1947), Skrjabin and Koval (1957) placed it under the family Fellodistomidae. In the present study latter arrangement is accepted.

This species resembles P. africana Fischthal and William (1971) in the shape of the body and in having longer hindbody and posttesticular space but differs from it in the absence of post-oral ring, in the shape and size of cirrus sac, which is small instead of long and retort-shaped, position of genital pore, which is on the anterior margin of acetabulum instead of posterior part of pharynx, size of the testes, which are much smaller than ovary,

Ovary pretesticular and intertesticular, submedian, diagonally oval, measures 0.17 x 0.010. Receptaculum seminis large. Uterus very extensive, ascending and descending coils extend from pharynx to posterior end of the body. Eggs numerous, oval, yellow, small. Vitelline follicles lateral, extending from oral sucker to posterior end of the body, confluent in posttesticular region.

Discussion

Linton (1911) proposed the name pyncadena to replace the generic name Didymorchis erected by him in 1910, since it was preoccupied. Srivastava, C.B. (1962) described Pyncadena komiyai and placed it under the family Allocreadiidae following Yamaguti (1958), Manter (1947), Skrjabin and Koval (1957) placed it under the family Fellodistomidae. In the present study latter arrangement is accepted.

This species resembles P. africana Fischthal and William (1971) in the shape of the body and in having longer hindbody and posttesticular space but differs from it in the absence of post-oral ring, in the shape and size of cirrus sac, which is small instead of long and retort-shaped, position of genital pore, which is on the anterior margin of acetabulum instead of posterior part of pharynx, size of the testes, which are much smaller than ovary,

1. Post-oral muscular ring present 2
Post-oral muscular ring absent 4

2. Body-broad, oval, nearly as broad as long or breadth equal to $3/4$ length, gonads situated near posterior extremity. 3

Body oval, hind body larger, gonads in anterior half of hind body

P. africans

3. Cirrus pouch short, lying at the anterior border of acetabulum, genital pore median at the base of pharynx, vitellaria diffused extending from pharynx to hind end of the body

P. lata

Cirrus pouch long, slender, genital pore submedian, near left margin of body, at level of anterior margin of pharynx, vitellaria very limited extending from anterior margin of acetabulum to ends of intestinal caeca

P. pyriformae

4. Testes symmetrical, uterus extending up to posterior end of the body, eggs with oculated embryos

P. bariliusi

Testes tandem, uterus extending up to testes, eggs not embryonated parasites of fresh water fishes

P. komiyai

- - - - -

- | | | |
|----|--|----------------------|
| 1. | Post-oral muscular ring present | 2 |
| | Post-oral muscular ring absent | 4 |
| 2. | Body-broad, oval, nearly as broad as long or breadth equal to 3/4 length, gonads situated near posterior extremity. | 3 |
| | Body oval, hind body larger, gonads in anterior half of hind body | <u>P. africans</u> |
| 3. | Cirrus pouch short, lying at the anterior border of acetabulum, genital pore median at the base of pharynx, vitellaria diffused extending from pharynx to hind end of the body | <u>P. lata</u> |
| | Cirrus pouch long, slender, genital pore submedian, near left margin of body, at level of anterior margin of pharynx, vitellaria very limited extending from anterior margin of acetabulum to ends of intestinal caeca | <u>P. pyriformae</u> |
| 4. | Testes symmetrical, uterus extending up to posterior end of the body, eggs with occulated embryos | <u>P. bariliusi</u> |
| | Testes tandem, uterus extending up to testes, eggs not embryonated parasites of fresh water fishes | <u>P. komiyai</u> |

- - - - -

Testes obliquely tandem, median, spherical or oval, postacetabular, intercaecal. Anterior testis smaller than posterior testis and measures $0.03 - 0.24 \times 0.05 - 0.24$. Posterior testis measures $0.078 - 0.25 \times 0.05 - 0.24$. Cirrus sac oval to club shaped, situated in between intestinal bifurcation and acetabulum, measures $0.08 - 0.18 \times 0.03 - 0.11$. Vesicula seminalis bipartite.

Ovary small, oval, submedian or median, attached to acetabulum, measures, $0.05 - 0.17 \times 0.04 - 0.10$. Receptaculum seminis and laurer's canal present. Shellgland complex large, uterus pretesticular. Vitellaria consists of large number of follicles extending from posterior margin of acetabulum and reaching almost upto hind end of the body and almost confluent in posttesticular region.

Excretory bladder tubular extending upto posterior border of posterior testis. Eggs large, yellow, oval embryonated, filamented and measures $0.07 - 0.10 \times 0.03 - 0.05$.

Discussion

This species has been described by Pande (1937) from the intestine of Channa punctatus from Handia, Allahabad and Haldwani. Kaw (1950) recorded this species from some other vertebrates, Coil and Kuntz, 1960 described it from small intestine of Channa punctatus from Dacca (Bangla Desh).

Testes obliquely tandem, median, spherical or oval, postacetabular, intercaecal. Anterior testis smaller than posterior testis and measures $0.03 - 0.24 \times 0.05 - 0.24$. Posterior testis measures $0.078 - 0.25 \times 0.05 - 0.24$. Cirrus sac oval to club shaped, situated in between intestinal bifurcation and acetabulum, measures $0.08 - 0.18 \times 0.03 - 0.11$. Vesicula seminalis bipartite.

Ovary small, oval, submedian or median, attached to acetabulum, measures, $0.05 - 0.17 \times 0.04 - 0.10$. Receptaculum seminis and laurer's canal present. Shellgland complex large, uterus pretesticular. Vitellaria consists of large number of follicles extending from posterior margin of acetabulum and reaching almost upto hind end of the body and almost confluent in posttesticular region.

Excretory bladder tubular extending upto posterior border of posterior testis. Eggs large, yellow, oval embryonated, filamented and measures $0.07 - 0.10 \times 0.03 - 0.05$.

Discussion

This species has been described by Pande (1937) from the intestine of Channa punctatus from Handia, Allahabad and Haldwani. Kaw (1950) recorded this species from some other vertebrates, Coil and Kuntz, 1960 described it from small intestine of Channa punctatus from Dacca (Bangla Desh).

Host : Clarias batrachus (Linn.)
Heteropneustes fossilis (B1.)
Channa striatus (B1.)
Channa punctatus (B1.)

Location : Intestine

Locality : Lalitpur, (U.P)

- - - - -

Family : ALLOCREADIIDAE Stossich, 1903
Sub family : ORIENTOCREADIINAE Yamaguti, 1958
Genus : Orientocreadium Tubanqui, 1931

Orientocreadium batrachoides Tubanqui, 1931
(Fig. 3)

Two samples of Orientocreadium batrachoides were collected from the intestine of Clarias batrachus (Linn.) during the two year study period from a pond at Moth in District Jhansi.

Description

The body is spinose, elongated, spindle shaped, with rounded anterior and blunt posterior end, measures 1.586×0.32 . The maximum breadth is at the level of anterior testis. Oral sucker is subterminal, rounded and measures 0.14×0.15 . Acetabulum is submedian, rounded and measures 0.14×0.15 roughly equal to the oral sucker. Prepharynx is large, measures 0.017×0.045 . Pharynx is well developed, measures 0.078×0.082 . Oesophagus is very small. Intestinal caecae are simple, long, terminating at posterior extremity.

The testes are rounded, entire, sub-equal, tandem, postequatorial, measuring $0.13 - 0.15 \times 0.28 - 0.25$. Cirrus sac is long, curved, lying lateral to acetabulum and

extends well behind it. It contains a vesicula seminalis interna, pars-prostatica and cirrus. Vesicula seminalis externa is long, saccular. Male and female genital opening are separate, preacetabular and median.

Ovary is oval, pretesticular, equatorial, intercaecal, lying between acetabulum and anterior testis and measures 0.12×0.26 . Receptaculum seminis is present. Vitelline follicles extend from the level of ovary upto hind end of the body. In posttesticular region, the follicles of two sides merge together. Eggs numerous, oval and measures 0.018×0.011 .

Excretory bladder is tubular with terminal excretory pore.

Discussion

The genus Orientocreadium was established by Tubangui, in 1931 to include Orientocreadium batrachoides, parasitic in Clarias batrachus (Linn.). In 1934, Yamaguti added Orientocreadium indicum from Rita buechanani and Orientocreadium pseudobagri from Pseudobagrus aurantiacus. He considered the genera Ganada Chatterji, 1933; Neoganada Dayal, 1938 and Nizamia Dayal, 1938 as synonymous with the genus Orientocreadium and also transferred their species under it.

The type species Orientocreadium batrachoides Tubangui, has been described in detail by many workers with little variations. These include Beverley-Burton, (1962) from Clarias mossambicus and Clarias mellandia; Fischthal and Kuntz, 1963; Kakaji, 1969 from Rita rita; Pandey, 1971 from Ophiocephalus punctatus and Jain and Chandra (1977) from Channa punctatus.

Yamaguti (1958) has described twelve species under the genus Orientocreadium. Orientocreadium ottoi has been considered a synonym of O. batrachoides Tubangui (1931).

This species has not been described from this locality, it is recorded herein.

Host : Clarias batrachus (Linn.)
Location : Intestine
Locality : Moth, District Jhansi, (U.P).

- - - - -

Family : ALLOCREADIIDAE Stossich, 1903
Sub family : ORIENTOCREADIINAE Yamaguti, 1958
Genus : Orientocreadium Tubangui, 1931

Orientocreadium indicum Pande, 1934

(Fig. 4)

The author collected 127 specimens of a trematode belonging to the genus Orientocreadium Tubangui, 1931 from the intestine of Clarias batrachus (Linn.). In all 360 specimens of the said host were examined of which 21 were found infected. This parasite was also recovered from the intestine of Heteropneustes fossilis (B1.) 360 specimens of this host were examined of which three were found infected and seven specimens were collected. All the three species of Channa - namely Channa punctatus (B1.) Channa striatus (B1.) and Channa marulius (Ham.) available locally were found heavily infected with this trematode. Thus out of 360 specimens of Channa punctatus examined, ten trematodes of this species were collected from three specimens. Out of 360 specimens of Channa striatus examined, three fishes were found infected and six trematodes were collected; out of 360 specimens of Channa marulius one was found infected with three specimens of this parasite.

On subsequent study it appears to be Orientocreadium indicum Pande, 1934 and has been described as such.

Description

Body small, elongated, spinose with rounded anterior and bluntly tapering posterior extremities, measuring $1.2 - 2.5 \times 0.24 - 0.34$, maximum width being in preequatorial region. Oral sucker spherical, subterminal, $0.08 - 0.13 \times 0.10 - 0.13$. Well developed prepharynx is present, $0.05 - 0.09$ in length, followed by muscular pharynx $0.08 - 0.10 \times 0.09 - 0.11$. Oesophagus very small. Intestinal caecae long, reaching upto posterior end. Acetabulum spherical, preequatorial, almost equal to oral sucker, measuring $0.13 - 0.19$.

Testes median, postequatorial with entire margin, measures $0.14 - 0.19 \times 0.09 \times 0.09 - 0.14$ and $0.15 - 0.21 \times 0.09 - 0.14$ respectively. Cirrus Sac large, spined, crescent shaped to the right of acetabulum, $0.24 - 0.32$ in length, enclosing pear shaped vesicula seminalis, pars prostatica, ejaculatory duct and protrucible cirrus. Vesicula seminalis externa in between ovary and acetabulum is present. Genital pore submedian, preacetabular.

Ovary median or submedian, postequatorial, situated midway between acetabulum and anterior testis, spherical,

0.09 - 0.12 x 0.06 - 0.10. Receptaculum seminis is absent, laurer's canal is present. Uterus extensive, occupying the whole postacetabular space, terminating in metraterm, opening in genital atrium. Vitellaria extend from the anterior border of ovary to posterior end of the body, where the follicles of the two sides meet. Eggs numerous, oval, operculate, yellow, 0.030 - 0.36 x 0.010 - 0.017.

Excretory bladder extending upto posterior margin of testes, excretory pore terminal.

Discussion

Pande (1934) described this new species Oriento-creadium indicum from the intestine of Rita buchanani from river Gomati at Jaunpur, Uttar Pradesh. This species differed from type species O. batrachoides Tubanqui, 1931 by the presence of spined cirrus and metraterm, spherical ovary instead of oval, maximum width in preequatorial region, vitellaria extending from posterior margin of acetabulum to posterior end of body where the follicles of two sides meet without forming lattice.

Yamaguti (1954, 1958), Saksena (1958, 1960), Gupta (1961) and Khalil (1961) accepted it as a valid species. Fischthal and Kuntz (1963) also accepted O. indicum as a valid species, characterised by the presence of spined

cirrus and metraterm, characters lacking in O.batrachoides and completely overlooked by Beverley-Burtan. Thus O.indicum is a valid species.

The present material forms new^host and locality record. For the first time the specimen has been recorded from a number of air-breathing fishes like Clarias batrachus (Linn.), Heteropneustes fossilis (B1.), Channa punctatus (B1.) Channa striatus (B1.) and Channa marulius (Ham.).

Host	:	<u>Clarias batrachus</u> (Linn.)
	:	<u>Heteropneustes fossilis</u> (B1.)
	:	<u>Channa punctatus</u> (B1.)
	:	<u>Channa striatus</u> (B1.)
	:	<u>Channa marulius</u> (Ham.)
Location	:	Intestine
Locality	:	Jhansi, (U.P.)

- - - - -

Family : ALLOCRAIDIIDAE Stossich, 1903
Sub family : ORIENTOCREADIINAE Yamaguti, 1958
Genus : Orientocreadium Tubangui, 1931

Orientocreadium pseudobagri Yamaguti, 1934

(Fig. 5)

The author collected 17 specimens of a trematode belonging to the genus Orientocreadium Tubangui, 1931 from the intestine of Clarias batrachus (Linn-). In all 360 specimens of the said host were procured from the river Betwa in District Hamirpur and examined, of which eight were found infected. This parasite was also recovered from the intestine of Heteropneustes fossilis (B1.) 360 specimens of this host were examined of which two were found infected and four parasites were collected this parasite was also recovered from the intestine of Channa punctatus (B1.) 360 specimens of this host were examined of which three were found infected and four trematodes were collected.

Description

Body spindle shaped, spinose with rounded anterior end and bluntly tapering posterior extremity, measures 2.8 - 3.10 x 0.53 - 0.55. Oral sucker is simple, subterminal oval, 0.12 - 0.14 x 0.14 - 0.17. Prepharynx is short. Pharynx comparatively large, oval, 0.038 - 0.09 x 0.13 - 0.14, Oesophagus long, measures 0.33 - 0.39. Intestinal

caecae long and reach nearly upto posterior extremity of body. Acetabulum spherical, preequatorial, 0.16 - 0.18.

Testes postequatorial, entire, oval or spherical, subequal or equal, tandem and measuring 0.17 - 0.20 x 0.19 - 0.21 and 0.18 - 0.21 x 0.18 - 0.20. The vesicula seminalis externa and vesicula seminalis interna are present. Vesicula seminalis externa occupies expanded basal part of the cirrus pouch and is elongated and oval in shape; vesicula seminalis interna is short, retort shaped, lies in front of ovary. The cirrus pouch is crescent shaped, running close to acetabulum on its right side, opening into genital atrium. Genital pore is preacetabular submedian and postbifurcal.

Ovary is pretesticular, round to oval, median, lies between acetabulum and anterior testis and measures 0.16 - 0.17. Shellgland complex behind the ovary. Uterus extends upto hind end of the body. Eggs numerous, oval, yellow, 0.020 - 0.026 x 0.005 - 0.008. Vitelline follicles extend from the mid acetabular region upto the hind end of body beyond the posttesticular region.

Excretory bladder is wide with terminal excretory pore.

Discussion

Tubangui (1931) created the genus Orientocreadium

and placed it under the family Allocreadiidae and later (1933) under the subfamily Allocreadiinae. McMullen (1937) erected the family Macroderoidae. Which was accepted by La Rue (1957) and Mehra (1966).

Yamaguti (1958) created the subfamily Orientocreadiinae to include the genera Orientocreadium and Macrotrema under the family Allocreadiidae. In the present work the family Allocreadiidae has been accepted.

Yamaguti (1958) has described 12 species under the genus Orientocreadium. The present specimens compare well with Orientocreadium pseudobagri Yamaguti, 1934, except some minor individual variations like oval oral sucker, small prepharynx, long oesophagus and extension of vitellaria halfway between thecaudal testis and posterior tip of the body and cirrus pouch on the right side of acetabulum instead of overlapping it. These variations may be considered as individual variations, so the present specimens are referred to as O. pseudobagri Yamaguti, 1934.

This is the first ^{new} locality record of this species.

Host	: <u>Channa punctatus</u> (B1.)
	: <u>Heteropneustes fossilis</u> (B1.)
	: <u>Clarias batrachus</u> (Linn.)
Location	: Intestine
Locality	: Hamirpur, (U.P.)

- - - - -

Family : ALLOCREADIIDAE Stossich, 1903
 Sub family : ORIENTOCREADIINAE Yamaguti, 1958
 Genus : Orientocreadium Tubangui, 1931.

Orientocreadium keni n.sp.

(Fig. 6)

Fifteen worms of this species were collected from the intestine of Clarias batrachus (Linn.) and Heteropnoustes fossilis (Bl.) from the river Ken, in District Banda (U.P.).

Description

Body spinose, small, elongated, measures 1.6 - 1.9 x 0.27 - 0.37. Maximum width being in pre-equatorial region. Oral sucker sub-terminal, rounded, 0.10 - 0.13 x 0.12 - 0.13. Prepharynx present, measures 0.02 - 0.05 in length. Pharynx spherical, muscular, measures 0.09 - 0.11 x 0.010 - 0.12. Oesophagus very small or absent. Intestinal caecae terminate at posterior extremity. Acetabulum spherical, pre-equatorial, 0.14 - 0.18.

Testes sub-median, transversely elongated, tandem, postequatorial, oval with entire margin, measures 0.12 - 0.17 x 0.07 - 0.12 and 0.13 - 0.19 x 0.07 - 0.12 respectively. Cirrus sac large, crescent shaped, 0.22 - 0.33 in length, situated to the right of the acetabulum, enclosing pear shaped vesicula seminalis, well developed pars prostatica, long, spined ejaculatory duct and protrucible cirrus. Well

developed coiled vesicula seminalis externa situated in between acetabulum and ovary. Genital pore preacetabular, submedian.

Ovary median, postequatorial, transversely elongated, situated midway between acetabulum and anterior testis, oval, $0.08 - 0.11 \times 0.06 - 0.09$. Shellgland complex posterolateral to ovary. Receptaculum seminis absent. The Laurer's canal present. Uterus very extensive, occupying the whole postacetabular space with descending and ascending coils. Vitellaria consists of large number of follicles, extending from posterior border of acetabulum to posterior end of the body. Eggs numerous, yellow, oval, operculated, $0.023 - 0.034 \times 0.011 - 0.016$ ^{m.m.} in size.

Discussion

The present species combined the characters of Orientocreadium indicum as well as O. pseudobagri but has its own distinctive features. It differs from both in having small body, transversely elongated ovary and testes, very small oesophagus, eggs rounded or oval.

Thus the present form differs from all other species it is therefore regarded as a new species and thus a new form Orientocreadium srivastavi has been added by the author in the present work.

This new form has been named after the name of the river Ken, from where these worms were recovered.

Host : Clarias batrachus (Linn.)
: Heteropneustes fossilis (Bl.)

Location : Intestine

Locality : Banda, (U.P.).

- - - - -

Family : GORGODERIDAE Loess, 1901
Sub family : PHYLLODISTOMINAE Yamaguti, 1958
Genus : Phyllodistomum Braun, 1899

Phyllodistomum tripathi Motwani & Srivastava, 1961
(Fig. 7)

Twelve specimens of Phyllodistomum tripathi Motwani and Srivastava, 1961 were collected from the intestine of Heteropneustes fossilis (B1.) during the two year study period from Barwa dam, in District Banda (U.P.).

Description

The body is aspinose, flask shaped, divisible into a narrow, tubular fore body and a foliate hind body with wavy margin and measures 1.63 x 2.31 mm in length and maximum width 0.75 x 1.21 mm just behind the posterior margin of acetabulum. Oral sucker is terminal, rounded, measures 0.208 - 0.320 x 0.210 - 0.318 in size. Acetabulum is larger than oral sucker, intercaecal, circular and measures 0.298 - 0.380 x 0.242 - 0.380 in size. Prepharynx and pharynx are absent. Mouth directly leads into oesophagus. Oesophagus is tubular, measures 0.149 - 0.715 in length and bifurcates into two simple, unbranched intestinal caeca, which extend upto hind region of body.

The testes are two in number, deeply lobed, intercaecal, postequatorial, obliquely tandem, placed in the expanded part of the body. They are more or less equal and measures 0.216×0.242 in size. Cirrus Sac is absent. Seminal vesicle is saccular, postbifurcal, intercaecal and anterior to acetabulum. Testes are separated by uterine coils.

The ovary is pretesticular, postacetabular, intercaecal, lobed, overlaps the right vitellaria and measures $0.098 - 0.210 \times 0.136 - 0.325$ in size. Receptaculum seminis is absent. Uterus with descending and ascending limbs, occupy intertesticular, posttesticular area, intercaecal and extracaecal. Its terminal part forms a muscular metraterm. The genital pore is situated behind the intestinal bifurcation, pre-acetabular, intercaecal, post bifurcal, preequatorial. The eggs are numerous, oval, operculated, both embryonated and unembryonated measures $0.014 - 0.016$ mm in size.

Vitellaria consist of two lobed glands, lying behind the ventral sucker on each side of the body.

The excretory bladder is sigmoid with terminal excretory pore. Excretory pore is near posterior end of the body.

Discussion

Genus Phyllodistomum was erected by Braun (1899) with Phyllodistomum folium (Olfers, 1816) Braun, 1899 as the type. The parasites of this genus normally inhabit the urinary bladder and also the intestine of fishes. It is observed that the forms recorded from urinary bladder are invariably larger.

A total of 15 species (13 from fishes and 2 from amphibians) of the genus Phyllodistomum have been reported from India. A complete list has been given by Thomas (1958) and a key to Indian species by Gupta (1953). Rai (1971) has critically reviewed the Indian species of the genus Phyllodistomum and according to him the various species can be divided into three groups on the basis of relative size of oral sucker and acetabulum. The relative size of suckers oral and ventral is a specific character and is least variable in the species. Accordingly i.e. equal size of both suckers - P. loossi Kaw, 1950; acetabulum smaller than oral sucker - P. vachius Dayal, 1949; P. vittatusi Gupta, 1953; P. chauhani Motwani and Srivastava, 1969 and acetabulum larger than oral sucker - P. simili Nybelin, 1926; P. lewisi H.D. Srivastava, 1938; P. singhiai Gupta, 1951; P. tripathi Motwani and C.B. Srivastava, 1961 are described as valid species.

The present specimen in the collection of the author has acetabulum distinctly larger than oral sucker, P. simili Nybelin, 1926; P. tripathi Motwani and Srivastava 1961; P. folium Braun, 1899 exhibit this relative structure of suckers. The authors are in agreement with Kakaji, 1969 in considering P. tripathi Motwani and Srivastava, 1961 as a synonym of P. folium, as the presence or absence of notch on the posterior end of the body and relative size of various organs except the suckers are variable characters.

To the best of my knowledge the present form in the collection of author is Phyllodistomum tripathi Motwani and Srivastava, 1961 with wavy margins having thick folds along the margins. These characters can be regarded as variable characters.

Host	: <u>Heteropneustes fossilis</u> (B1.)
Location	: Intestine
Locality	: Banda (U.P.).

- - - - -

Family : ISOPARORCHIIDAE Poche, 1926
 Genus : Isoparorchis Southwell, 1913
 (Syn. Leptolecithum Kobayashi, 1915)

Isoparorchis hypselobagri (Billet, 1898) Odhner, 1911
 (Fig. 8)

Metacercariae of this species were collected from the body cavity of Channa punctatus (B1.), Channa striatus (B1.) and Channa marulius (Ham.) during the months of April, January and September from the river Ken, in District Banda (U.P.).

Description

Body large, cylindrical, unspined with thick cuticle, flesh coloured, translucent when expanded, foliate with bluntly projected anterior and broadly rounded posterior extremity and measures 1.02-4.05 x 0.62-1.07. Oral sucker is subterminal, oval, 0.16-0.39 x 0.17-0.49. Prepharynx absent, pharynx well developed. Oesophagus indistinguishable. Intestinal caeca long, serpentine, extending to near posterior end of the body. Acetabulum is larger than oral sucker, rounded, situated in the anterior half of the body and measures 0.27-1.09 x 0.27-1.2. Gonads indistinguishable. Genital pore situated between the two suckers at a distance of 0.31-0.61.

Excretory vesicle Y-shaped, excretory pore terminal.

Discussion

Isoparorchis hypselobagri (Billet, 1898) has been described as the type species of the genus Isoparorchis Southwell, 1913. The metacercaria of this species has been recorded from a number of fresh water fishes like Wallago attu and Barbus tor as has been listed by Pandey (1970) and Srivastava (1972). Srivastava (1977) discussed its host, distribution and relationships.

The writer has collected this specimen from all the species of Channa available locally, viz. Channa punctatus (B1.) Channa striatus (B1.) and Channa marulius (Ham.). This is the first host and locality record of the larva from Banda.

Host	:	<u>Channa punctatus</u> (B1.)
	:	<u>Channa striatus</u> (B1.)
	:	<u>Channa marulius</u> (Ham.)
Location	:	Body cavity
Locality	:	Banda (U.P.)

- - - - -

Family : HEMIURIDAE Luhe, 1901
 Sub family : HALIPEGINAE Ejsmont, 1931
 Genus : Genarchopsis Ozaki, 1925
 (Syn. Progonus looss, 1899 preoccupied)

Genarchopsis piscicola Srivastava, 1933.

(Fig. 9)

The present study is based on eight specimens of this species collected from intestine of Channa punctatus (B1.). 360 specimens of Channa punctatus (B1.) were procured from Dongri dam and Khaprar dam, in District Jhansi and dissected, of which intestine of four fishes were found infected with Genarchopsis.

Description

The body is small, fusiform, aspinose, muscular and measures 1.26-2.62 x 0.47-0.89. The oral sucker is subterminal, cuplike and measures 0.32-0.40 x 0.32-0.50. Acetabulum is large, postequatorial, well developed, muscular, almost spherical, measuring 0.60-0.87 x 0.61-0.87. The prepharynx is absent. Pharynx present, small and measures 0.11-0.13 x 0.02-0.19. Oesophagus absent. Intestinal caecae long and sinuous, extending upto posterior margin of body where they unite.

Testes oval, Postacetabular, slightly obliquely tandem, extracaecal and anterior testis measures 0.24-0.37 x 0.17-0.28 and posterior testis 0.24-0.37 x 0.17-0.33 respectively. The cirrus sac absent. Pars prostatica is tubular and convoluted. The ejaculatory duct is short and opens along with metraterm into a short hermaphroditic duct.

Ovary small, postacetabular, intercaecal, spherical, situated almost at the level of posterior testis and measuring 0.15-0.23 x 0.12-0.33. The uterine coils are intercaecal, extending posteriorly upto the hind border of the testes. The left vitelline follicles measure 0.12 - 0.33 x 0.9 - 0.42, while right follicles measure 0.18-0.52 x 0.07-0.28. The genital pore just below pharynx near the intestinal bifurcation.

Discussion

Looss (1899) erected the genus Progonus for Mulleri Levinsen, 1881. He (1902) renamed it as Genarches thinking the name Progonus as preoccupied by the insect genus Progona Berg, 1886. Führmann (1904, 1928), Odhner (1905) considered Progonus as valid. Ozaki (1925) proposed a new genus Genarchopsis to accommodate his new species G. goppo Srivastava, H.D. (1933) while accepting the name progonus valid, considered the genus Genarchopsis as synonym of the former. He also erected a new genus Ophiocorchis to accommodate his new species O. lobatum. Yamaguti (1958,

1971) considered the genera Progonus, Genarches, Ophiocorchis as congeneric with Genarchopsis, the former two being, in his opinion, preoccupied. This arrangement is more or less being followed by the subsequent workers. Rai (1971) made a detailed study of various Indian species related to the genus Ophiocorchis and Genarches and came to the conclusion that all Indian species are synonym of G. goppo Ozaki, 1925. He however, did not deal with the question of the validity of the genus Progonus. Srivastava, H.D. and Sahai (1978) tried to revalidate the genus Ophiocorchis. The genus Progonus is not preoccupied as generally believed. According to Article 56 of International Rules of Zoological Nomenclature, the genera Genarchopsis, and Ophiocorchis have therefore been related to the genus Progonus.

The trematode under discussion has been compared with the valid species of genus Genarchopsis Ozaki, 1925 namely G. ovocaudatum, G. piscicola and G. dasus. It differs from G. ovocaudatum in the length and maximum width of the body, size of ovary, ratio of suckers and extension of uterine coils. It also differed from G. dasus (Gupta, 1951) in the length and maximum width of body, length and width of oral sucker, ratio of suckers and measurement of *versicula seminalis*.

However, the worm approaches more closely to G. piscicola Srivastava, 1933 in the extension of uterine coils

and principal body measurement, except for the differences in the length of the body, the ratio of suckers and the nature of the uterine coils. These variations are minor ones and do not suggest the proposition of a new species and so the present worms are referred to as Genarchopsis Piscicola Srivastava, 1933.

Host : Channa punctatus (B1.)

Location : Intestine

Locality : Jhansi (U.P.).

- - - - -

Family : HEMIURIDAE Luhe, 1901
 Sub family : HALIPEGINAE Ejsmont, 1931
 Genus : Genarchopsis Ozaki, 1925
 (Syn. Progonus Looss, 1899
 preoccupied).

Genarchopsis goppo Srivastava, 1933
 (Fig. 10)

Out of 360 specimens of Channa punctatus (B1.) examined, the intestine of one was found infected with two worms of genus Genarchopsis Ozaki, 1925. The fishes were procured from Gunta dam at Mau, in District Banda, (U.P.).

Description

Body elliptical with both extremities rounded and measuring 3.6 - 4.89 x 1.07 - 1.9. Maximum breadth in acetabular zone. Oral sucker oval, subterminal, 0.32 - 0.36 x 0.39 - 0.44. Pharynx oval, 0.17 - 0.4 x 0.17 - 0.20. Acetabulum large, spherical, equatorial, 0.79 - 0.85 x 0.79 - 0.87.

Testes oval, postacetabular, almost at the same level right measuring 0.25 - 0.31 x 0.45 - 0.47 and left 0.35 - 0.36 x 0.44 - 0.45 respectively. The cirrus sac absent. Vesicula seminalis well developed, long, cylindrical, postbifurcal, lying free in parenchyma which continues

into oval, compact pars prostatica surrounded by prostate gland cells. Terminal part of pars prostatica joins with metraterm to form hermaphroditic duct. Genital pore submedian in level with pharynx.

Ovary oval, posttesticular, $0.16 - 0.30 \times 0.4 - 0.38$ lying just below right testis. Shellgland complex, postovarian. Receptaculum seminis uterinum present. Metraterm well developed, muscular, receiving pars prostatica at its distal end. Eggs with filament on one side. Vitellaria two lobed glands partly overlapping the intestinal caeca, lying in hind part of body.

Excretory bladder Y-shaped with the arms anastomosing dorsal to oral sucker.

Discussion

Srivastava (1933) described the type species Progonus lobata (Srivastava, 1933) from the stomach of Channa striatus from Lucknow. Gupta (1951) described Genarchopsis farugis from the intestine of Mastacembelus armatus which is considered as synonym of this species in the light of the variations observed. G. malanosticus Dwivedi, 1965 and G. cuchiai Kakji, 1969 are considered conspecific with P. lobata as the characters used by them to differentiate their new species fall within the limit of variations.

The present specimen, therefore, has been described as type species Genarchopsis goppo Srivastava, 1933. This is the first locality record of this specimen from Banda.

Host : Channa punctatus (B1.)
Location : Intestine
Locality : Mau (District Banda, U.P.)

- - - - -

Family : HEMIURIDAE Luhe, 1901
 Sub family : HALIPEGINAE Ejsmont, 1931
 Genus : Genarchopsis Ozaki, 1925
 (Syn. Progonus Looss, 1899
 preoccupied).

Genarchopsis singularis Srivastava, 1933
 (Fig. 11)

360 specimens of Channa punctatus (B1.) and 360 specimens of Channa striatus (B1.) were examined during present study. Out of these, four and one specimen were found infected respectively with Genarchopsis singularis Srivastava, 1933. In all ten worms were collected from the intestine of hosts. The fishes were procured from Rajghat dam, in District Lalitpur (U.P.).

Description

Body small, oval, 0.98 - 3.32 x 0.49 - 0.86 with maximum breadth at acetabular zone. Oral sucker oval, subterminal, 0.13 - 0.33 x 0.19 - 0.37. Pharynx oval, 0.08 - 0.2 x 0.07 - 0.10. Oesophagus small.

Testes symmetrical, postacetabular, oval, overlapping intestinal caeca, measuring 0.11 - 0.21 x 0.09 - 0.13 and 0.11 - 0.20 x 0.08 - 0.17 respectively. Cirrus sac absent.

Ovary small, spherical, median, posttesticular, lying in the hind part of body, anterior to vitelline follicles, $0.067 - 0.22 \times 0.087 - 0.16$. Shellgland complex postovarian. Uterus with transverse coils extending posteriorly upto vitelline zone which continues anteriorly as metraterm and opens into pars prostatica. Eggs yellowish oval, $0.021 - 0.037 \times 0.011 - 0.015$ with a filament on one side. Vitellaria two, compact, oval glands one on either side in posterior most part of the body.

Excretory bladder Y-shaped with arms anastomosing dorsal to oral sucker.

Discussion

Chauhan, 1953 in the comprehensive work on the family Hemiuridae maintained the genera Progonus and Ophiocorchis synonym of Genarches and transferred their species under it. Yamaguti, 1958 synonymised the genus Ophiocorchis Srivastava, 1933 (Progonus Looss, 1899, Preoccupied; Genarches Looss, 1902 Preoccupied) with Genarchopsis Ozaki, 1925.

Srivastava (1933) described Ophiocorchis singularis from the intestine of Channa marulius at Allahabad. Gupta (1951) collected some specimens from Channa punctatus at Lucknow and Saharanpur, U.P. and referred them to as new species Ophiocorchis indicum. Chauhan (1954) remarked that

the differences enumerated by Gupta do not merit the establishment of a new species and the two species should be considered identical.

The present specimens have been collected from Channa punctatus and Channa striatus from Rajghat dam in District Lalipur (U.P) and combine the characters of both the species. Thus it is evident that O. indicum is conspecific with O. singularis as suspected by Chauhan (1954). The present collection, however forms a new locality record.

Host : Channa punctatus (B1.)
 : Channa striatus (B1.)
 Location : Intestine
 Locality : Lalitpur, (U.P.)

Key to the species of the genus Progonus Looss, 1899 :-

- | | | |
|----|---|----------------------|
| 1. | Oesophageal pouch present
Vitellaria lobed or compact | 2 |
| | Oesophageal pouch absent,
Vitellaria compact | 4 |
| 2. | Cirrus pouch present | <u>P. thapari</u> |
| | Cirrus pouch absent | 3 |
| 3. | Vitelline glands distinctly lobed
acetabulum equatorial or post-
equatorial | <u>P. lobata</u> |
| | Vitelline glands compact
or crenulated | <u>P. singularis</u> |

4. Vitellaria a little anterior to
posterior extremity and uterus
extending beyond vitellaria P. mulleri
- Vitellaria near posterior
extremity, Uterus extending upto
vitellaria or still
anterior to it 5
5. Main mass of uterine loops
anterior to acetabulum P. piscicola
- Uterus equally anterior and
posterior to acetabulum 6
6. Uterus mostly intercaecal P. goppo
- Uterus extending into
extracaecal zone P. angullae
- - - - -

Family : CLINOSTOMIDAE Luhe, 1901

Genus : Clinostomum Leidy, 1856

Clinostomum complanatum (Rudolphi, 1819) Braun, 1899
(Metacercaria)

The author collected 40 specimens of Clinostomum complanatum (Rudolphi, 1819) Braun, 1899 throughout the year 1993 to 1995 from the body cavity of Channa punctatus (B1.).

The parasite resembles with the description given by previous workers and hence no description and measurement have been recorded.

Host : Channa punctatus (B1.)

Location : Body Cavity

Locality : Lalitpur (U.P.).

- - - - -

Family : CLINOSTOMIDAE Luhe, 1901
 Sub family : EUCLINOSTOMINAE Yamaguti, 1958
 Genus : Euclinostomum Travassos, 1928

Euclinostomum heterostomum (Rudolphi, 1809)
 Travassos, 1928
 (Fig. 12)

Metacercaria (Larval trematodes) of this species were collected in cysts from Body cavity of Channa punctatus (B1.). Out of 360 specimens of Channa punctatus (B1.) examined during 1993 to 1995, three fishes were found infected with these larval forms in the month of July and January. The fishes were procured from Rajghat dam, in District Lalitpur (U.P.).

Description

Body large, oval, unspined with rounded extremities measures 4.9 - 5.7 x 2.42 - 3.03 with collar like formation at anterior end. Oral sucker very small, oval, subterminal measures 0.3 - 0.44 x 0.3 - 0.36. Pharynx small, thick walled, measures 0.13 - 0.16 x 0.16 - 0.18, surrounded by loosely arranged parenchymal muscles extending from posterior border of oral sucker to the base of collar, roughly halfway between intestinal bifurcation and acetabulum. Larval eye-spots present on either side of Pharynx. Oesophagus absent. Intestinal caeca very thin upto posterior

border of acetabulum, after which lateral diverticulae start. Lateral diverticulae are single and branched terminally postero-lateral, extending upto lateral margins of the body their number being 11 on the left side and 12 on the right. Acetabulum very large, spherical, median, preequatorial, measuring $1.27-1.5 \times 1.2-1.29$ at a distance of $1.17-1.35$ from anterior end.

Testis postequatorial, tandem, median, intercaecal. Anterior testis crescent shaped and posterior testis y or V shaped, measuring $0.33-0.34 \times 0.77-0.79$ and $0.54-0.55 \times 0.54-0.58$ respectively. Vas-efferentia of posterior testis running parallel to right caeca and joins with vas-efferentia of the anterior testis, then enters the cirrus sac as vas-deferens. Cirrus sac small, oval, in front of anterior testis, in between its arms, enclosing bipartite seminal vesicle, pars prostatica and cirrus. Genital pore median, at the level of anterior third of cirrus sac.

Ovary small, oval, submedian, intertesticular, intercaecal, measuring $0.12-0.17 \times 0.12-0.2$. Oviduct, short arising from posterior border of ovary. Shellgland complex large, diagonally placed on one side of ovary. Metraterm very short and opens into genital atrium. Vitelline follicles very small, immature, lateral extending from posterior margin of acetabulum upto posterior margin of

acetabulum upto posterior end of body and continuous in posttesticular region.

Discussion

Travassos (1928) created the genus Euclinostomum with Euclinostomum heterostomum as its type species; a generic diagnosis was not given. Yamaguti (1958) created the sub-family Euclinostominae with Euclinostomum as its only genus.

The species E. heterostomum was first described as Distoma heterostomum by Rudolphi (1809). The description was rather brief and general dealing with external features and was without an illustration. Braun (1900) presented the first detailed account of the morphology of adult E. heterostomum from herons, Ardea purpurea, A. cinerea and Nycticorax griseus. Monning (1926) reported three metacercariae of E. heterostomum from the muscles of an unnamed fish. Joyeux and Houdemer (1928) recorded adult E. heterostomum from egrets, Garzetta garzetta. Metacercariae were found by them in the muscles of the fish, Anabas scandens.

Srivastava (1950) found metacercariae of E. heterostomum from Channa punctatus embedded in the liver and attached to the kidneys or muscles of the coelomic wall.

Adult worms occurred naturally in the night heron,
Nycticorax nycticorax.

Euclinostomum indicum was described by Bhalerao (1942) from the body cavity of Channa punctatus. Agrawal (1959) collected adult of E. indicum from herons, Bubulcus ibis fed with the fish, Channa punctatus, Fischthal and Kuntz (1963) considered E. indicum (Bhalerao, 1942, metacercaria; Agrawal, 1959 adult) synonym of E. heterostomum.

About a dozen species have been described so far under the genus Euclinostomum, of which five are from our country viz. E. heterostomum, E. bhagvantami, E. Channai, E. hepatocaecum and E. indicum. The characters utilized by previous workers for differentiating the known species of Euclinostomum are all highly variable and which include, the body shape, the structure of prepharynx, pharynx, oesophagus, number and shape of caecal diverticula. Perusal of literature shows that E. heterostomum the type species the genus, enjoys a wide host range and occurs in different geographical locality throughout the world. Its metacercarias are found in fresh water fishes consequently wide morphological variations in the anatomy are natural in this species.

The present specimens agree with the description given by the Braun (1900) except in the presence of well developed pharynx and vitellaria, which are considered as individual variations of E. heterostomum.

Host	:	<u>Channa punctatus</u> (B1.)
Location	:	Body cavity
Locality	:	Lalitpur, (U.P.)

- - - - -

Family : HETEROPHYIDAE Odhner, 1914
 Sub family : HAPLORCHIINAE Looss, 1899
 Genus : Haplorchoides Chen, 1949

Haplorchoides seenghali Dayal and Gupta, 1954
 (Fig. 13)

One specimen of this species was collected from the intestine of Heteropneustes fossilis (B1.) out of 360 specimens of Heteropneustes fossilis (B1.) examined throughout the year 1993, 1994 and 1995. The fishes were procured from Barwa dam, in District Banda (U.P.).

Description

Body very small, oval with rounded anterior and posterior extremities, 0.60 x 0.24. Oral sucker subterminal spherical, 0.06 in diameter. Prepharynx 0.05 in length. Pharynx oval, well developed 0.015 x 0.011. Oesophagus 0.012 in length. Intestinal caecae terminating a little anterior to hind end of the body. Acetabulum absent.

Testis single, very large, oval, intercaecal, 0.09 x 0.12. Cirrus pouch absent. Seminal vesicle bipartite. Genital sac globular situated on left intestinal caecum.

Ovary small, median, pretesticular, spherical, 0.04 - 0.05, Uterus very extensive occupying the whole

postbifurcal body, containing large, yellow, oval, filamented eggs occupying entire posttesticular region. Vitelline follicles extending from hind end of ovary to hind end of testis.

Discussion

Chen (1949) established the genus Haplorchoides with H. cahirinus (Looss, 1896) as its type species.

The question of the validity of the genera Haplorchis Looss, 1899; Monorchotrema Nishigori, 1924 and Chen, 1949 have been discussed by several authors. Witenberg (1929 and 1930), Srivastava (1935), Chen (1936) and Dawes (1946) considered the synonymy of Monorchotrema to Haplorchis.

Gohar (1934) and Dayal (1935) considered Haplorchis and Monorchotrema as distinct genera. Srivastava (1935) splitted the genus Haplorchis into two subgenera; Haplorchis (Monorchotrema) and Haplorchis (Haplorchis). The subgenus Haplorchis (Monorchotrema) is characterized in having prepharynx shorter than oesophagus, testis and vitellaria caudal in position and with a rudimentary acetabulum, while the subgenus Haplorchis (Haplorchis) is characterized in having prepharynx longer than oesophagus, testes and vitellaria more anterior and the acetabulum being absent.

Chen (1936) considered the synonymy of the genera Haplorchis Looss, 1899 and Monorchotrema Nishigori, 1924 and considered Haplorchis to be the only valid genus and Monorchotrema to be identical and synonymous to it. In 1949, he differentiated the two genera Haplorchis Looss, 1899 and Haplorchoides n.g. on the basis of relative size of prepharynx and oesophagus, a short prepharynx, long oesophagus, thin-walled vesicula seminalis and more posterior position of testis and vitellaria, Characterized Haplorchis while a long prepharynx, short rudimentary oesophagus and more forward position of testis and vitellaria characterized Haplorchoides. Yamaguti, 1954 considered Pseudohaplorchis Dayal, 1949 as a synonym of Haplorchoides Chen, 1949.

Gupta (1953), Yamaguti (1958 and 1971) and Agrawal, 1964 accepted Chen's (1949) view and considered the genus Haplorchoides as distinct from Haplorchis whereas Chatterji (1953 and 1956) and Baugh (1963) considered Haplorchoides as a synonym of Haplorchis. Later Pearson (1964) accepted this genus and placed it under the family Heterophyidae on the basis of the position and shape of excretory bladder. Rai and Pande (1967) agreed with Pearson (1964) and placed it in the sub-family Haplorchiinae Looss, 1899 on the basis of the presence of armed acetabulum embedded inside the ventrogenital sac and saccular and post-testicular excretory bladder. In the present study, the latter arrangement has been accepted.

Haplorchoides seenghali has been described by Dayal and Gupta (1954) from the intestine of Mystus seenghala. Gupta (1955) gave complete description of the species but described it as a new species.

The present specimens resemble type specimen except for smaller size of body, in principal body measurements and very large size of testis, which may be regarded as variable characters. So the specimen is described as Haplorchoides seenghali Dayal and Gupta, 1954.

Host	: <u>Heteropneustes fossilis</u> (B1.)
Location	: Intestine
Locality	: Banda (U.P).

- - - - -

P A R T - I I I

HOST - PARASITE RELATIONSHIP

HOST-PARASITE RELATIONSHIP

Study of helminthological literature from India indicates that more attention has been paid to taxonomic studies of helminth worms. While the taxonomy forms the basis of all biological investigations, still it alone can not fulfill the objectives of parasitologists. Host-parasite relationship has to be given due weightage to assess the exact extent of damage caused by these parasites. Till now very little attention has been given to host-parasite relationship, population biology and estimation of helminth infection with regard to most of the species. These lacunae are mainly responsible for our failure in implementing our objectives and in combating helminth infections.

In order to make the faunistic studies more objective, host-parasite relationship has been studied and discussed in detail. Investigations on these relatively explored branches are bound to reveal important clues for a correct understanding of the nature and extent of their pathogenic role.

A parasite is always under the influence of two types of environments - viz; the internal environment in which the parasite lives and the external environment in which the host lives. It is the interaction of the influence of

these environments and the strategy adopted by the parasites to counter influence that develops the host specificity and host parasite relationship. Thus the establishment and survival of helminths in their hosts is controlled by the internal environment of the host as well as its external environment.

Thus the major environmental factor that influence the incidence of helminth infection and host-parasite relationship are both biotic and abiotic. In the present study various abiotic factors including - Index of total helminth infection in trematodes, host - wise analysis, overall incidence, level and intensity of parasitization, seasonal incidence have been studied and statistically analysed, similarly biotic factors like the influence of sex of the host and size of the host have also been studied.

The climate of Bundelkhand region is subtropical and four seasons which can be recognised are the Winter season (Nov., Dec., Jan., Feb.), Summer season (March, April, May, June) and Rainy season (July, Aug., Sept., Oct.).

The average atmospheric temperature ranges from 25.11°C (in Jan.) to 41.35°C (May) and minimum temperature ranges from 10.51°C (in Jan.) to 26.58°C (May). The temperature gradually shows a decreasing trend from July to December and increasing trend from January to May.

Relative humidity is more in morning (8.30 am) than in evening (5.30 pm). The minimum in morning hours was 24.90% in April while maximum 85.35% was in the month of August. In the evening the lowest humidity values were ^{in MAY} 16.50% and highest 85.33% in August.

Since during the present study period trematodes formed the predominant helminth parasites, hence the present study is mainly concentrated on the host-parasite relationship of trematodes. As the cestode recovered was only one, and no nematodes were reported, so they are not being statistically analysed.

Under the present project, the helminth parasites of five different host fishes viz. - Channa punctatus (B1.); Channa striatus (B1.); Channa marulius (Ham.); H. fossilis (B1.) and Clarias batrachus (Linn.) were collected from July 1993 to June 1994 and from July 1994 to June 1995. A total of 1800 host fishes were collected and examined regularly during that period. On an average fifteen fishes of each host species (Total about 75 fishes per month) were examined. The data thus collected has been statistically analysed.

INDEX OF TOTAL HELMINTH INFECTION (TREMATODES)

INDEX OF TOTAL HELMINTH INFECTION (TREMATODES)

A total of 1800 host fishes belonging to five host species were procured during two years period from July 1993 to June 1995. The data collected was analysed to determine percentage of total helminth infection.

TABLE I

Month & Year		No. of parasite recovered	Percentage of trematode infestation
July	93-94	28	18.66
August	93-94	70	46.66
September	93-94	29	19.33
October	93-94	26	17.33
November	93-94	27	18.00
December	93-94	24	16.00
January	94-95	38	25.33
February	94-95	28	18.66
March	94-95	19	12.66
April	94-95	16	01.66
May	94-95	01	00.66
June	94-95	02	01.33

It is clear from the above Table 1 that the highest % of trematode infestation was in the month of August and lowest in the month of May and June.

It was observed from separate study of year wise incidence, that the helminth infestation was relatively less in the year 1993-94 as compared to that in 1994-95 i.e. there is a definite increase in the rate of helminth infestation during 1994-95. (Plate 14).

From the study of monthly fluctuation of trematode parasites, it is evident that the infection shows a decline during summer months prior to breeding season (May - June) and increases in rainy season. The reason for this may be that prior to breeding season, the fishes stop feeding for some time and so there is very little chance of their getting infected, but during rainy season water gets polluted, so the infestation rate is high.

HOST - WISE ANALYSIS OF PARASITES

HOST-WISE ANALYSIS OF PARASITES

Host-wise analysis of various parasites recovered has been analysed in Table 2A, 2B, 2C, 2D & 2E respectively.

Channa punctatus (B1.)

TABLE 2-A

Parasite	Habitat	Total No. of parasites recorded during July 1993 to June 1995	Total No. of positive hosts	Month and Year in which infection was recorded
(a)	(b)	(c)	(d)	(e)
<u>Genarchopsis singularis</u> <u>Srivastava, 1933</u>	Intestine	8	4	July 1994 August 1994 December 1994 April 1995
<u>Genarchopsis goppo</u> <u>Srivastava, 1933</u>	Intestine	2	1	February 1995
<u>Genarchopsis piscicola</u> <u>Srivastava, 1933</u>	Intestine	8	4	September 1994 November 1994 December 1994 March 1995

(a)	(b)	(c)	(d)	(e)
<u>Orientocreadium indicum</u> Pande, 1934	Intestine	10	3	July August November 1993 1994 1994
<u>Orientocreadium pseudobagri</u> Yamaguti, 1934	Intestine	4	3	August August November 1993 1994 1994
<u>Allocreadium handiai</u> Yamaguti, 1934	Intestine	5	4	November December July April 1993 1993 1994 1995
Metacercaria				
<u>Euclinostomum heterostomum</u> (Rudolphi, 1809) Travassos, 1928	Body cavity	4	3	July January 1994 1995
<u>Isoparorchis hypselobagri</u> (Billet, 1898) Odhner, 1911)	Body cavity	3	1	April 1994
<u>Clinostomum complanatum</u> (Rudolphi, 1819) Braun 1899	Body cavity	40	17	July to Dec Jan to June 1993 1994

Channa striatus (B1.)

TABLE 2-B

<u>Parasite</u>	<u>Habitat</u>	<u>Total No. of parasites recor- ded during July 93 to June 1995</u>	<u>Total No. of posi- tive hosts</u>	<u>Month & Year in which infection was recorded</u>
<u>Genarchopsis singularis</u> <u>Srivastava, 1933</u>	Intestine	2	1	October 1994
<u>Orientocreadium indicum</u> <u>Pande, 1934</u>	Intestine	6	3	July 1994 September 1994 November 1994
<u>Allocreadium handiei</u> <u>Pande, 1937</u>	Intestine	3	2	August 1994 April 1995
<u>Metacercaria</u>				
<u>Isoparorchis hypselobagri</u> <u>(Billet, 1898)</u> <u>Odhner, 1911</u>	Body cavity	2	1	January 1995

Channa marulius (Ham.)

TABLE 2-C

Parasite	Habitat	Total No. of parasites recorded during July 93 to June 95	Total No. of positive hosts	Month & Year in which infection was recorded
<u>Orientocreadium indicum</u> Pande, 1934	Intestine	3	1	August 1994
<u>Metacercaria</u>				
<u>Isoparorchis hypselobagri</u> (Billet, 1898) Odhner, 1911	Body cavity	2	1	September 1994

Heteropneustes fossilis (B1.)

TABLE 2-D

Parasite	Habitat	Total No. of parasites recorded during July 93 to June 95	Total No. of positive hosts	Month & Year in which infection was recorded
<u>Orientocreadium indicum</u> Pande, 1934	Intestine	7	3	March 1994 July 1994 November 1994
<u>Orientocreadium pseudobagri</u> Yamaquti, 1934	Intestine	4	2	February 1994 February 1995
<u>Orientocreadium keni</u> n.sp.	Intestine	2	1	January 1995
<u>Allocreadium handiai</u> Pande, 1937	Intestine	5	1	February 1994
<u>Haplorchoides seenghali</u> Dayal and Gupta, 1954	Intestine	1	1	February 1995
<u>Phyllodistomum tripathi</u> Motwani and Srivastava, 1961	Intestine	12	5	July 1994 August 1994

TABLE 2-E

Parasite	Habitat	Total No. of parasites recorded during July 93 to June 95	Total No. of positive hosts	Month & Year in which infection was recorded
<u>Orientocreadium indicum</u> Pande, 1934	Intestine	127	21	Aug, Sep., Dec. 1993 Feb, Mar., Aug. Sep, Oct., Nov. 1994 December 1994 Jan, Feb., Mar April 1995
<u>Orientocreadium pseudobagri</u> <u>Yamaguti, 1934</u>	Intestine	10	8	Aug, Sep., Oct, 1994 November 1995 Jan, Feb., Mar.
<u>Orientocreadium keni</u> n.sp.	Intestine	13	7	Jan. Feb., Aug. October 1994 Feb., March 1995
<u>Orientocreadium batrachoides</u> <u>Tubangui, 1931</u>	Intestine	2	1	April 1995
<u>Allocreadium handiai</u> <u>Pande, 1937</u>	Intestine	15	8	November 1993 Mar, Apr., Oct. 1994 Feb. Mar., Apr. 1995
<u>Pycnadena barilii</u> <u>V. Kumari, 1973</u>	Stomach	1	1	January 1995

- Channa punctatus (Bl.). From July 1993 to June 1995, the author regularly examined fifteen fishes per month of each host species. Accordingly Channa punctatus was collected every month to procure parasites. On an average fifteen host fishes were examined per month and one hundred & eighty fishes were examined from July 1993 to June 1994 and the same number was examined in July 1994 to June 1995 (Total 360 host fishes). The number of trematodes and the month in which they were recovered were recorded. The analysis of the data has been given in (Table 2-A). This table shows that in all, Channa punctatus harboured nine trematode species including 3 metacercaria. Orientocreadium indicum was the dominant trematode species.
- Channa striatus (Bl.). Fifteen host fishes per month were examined for a period of two years. The data collected has been analysed in (Table 2-B). The analysis shows that C. striatus harboured four trematode species including one metacercaria. Parasite-wise, O. indicum formed most dominant trematode species.
- Channa marulius (Ham.). Channa marulius showed minimum trematode infestation during a study period of two years from July 1993 to June 1995. This species of Channa harboured only one trematode species O. indicum and one metacercaria.

- H.fossilis (B1.). This air-breathing host fish harboured six trematode species including one new species with maximum number of P.tripathi. The new species harboured by this host is O. keni.
- Clarias batrachus (Linn.). Most commonly available cat fish, Clarias batrachus, harboured six different trematode species including a new species Orientocreadium keni.

Thus the above analysis suggests that Channa punctatus (B1.) and Clarias batrachus (Linn.) are the most susceptible host fishes for trematode infestation.

- - - - -

OVERALL INCIDENCE OF TREMATODES

OVERALL INCIDENCE OF TREMATODES

Channa punctatus (B1.)

TABLE 3-A

Parasite	No. of indivi- duals parasi- tized	Parasiti- zation rate percen- tage	Mean No. of parasite per host.
<u>Genarchopsis singularis</u> Srivastava, 1933	4	1.11	2.00
<u>Genarchopsis goppo</u> Srivastava, 1933	1	0.277	2.00
<u>Genarchopsis piscicola</u> Srivastava, 1933	4	1.11	2.00
<u>Orientocreadium indicum</u> Pande, 1934	3	0.833	3.33
<u>Orientocreadium pseudobagri</u> Yamaguti, 1934	3	0.833	1.33
<u>Allocreadium handiai</u> Pande, 1937	4	1.11	1.25
Metacercaria			
<u>Euclinostomum heterostomum</u> (Rudolphi, 1809) Travassos, 1928	3	0.833	1.33
<u>Isoparorchis hypselobagri</u> (Billet, 1898) Odhner, 1911	1	0.277	3.00
<u>Clinostomum complanatum</u> (Rudolphi, 1819) Braun 1899	17	4.722	2.352

Channa striatus (B1.)TABLE 3-B

Parasite	No. of indivi- duals parasi- tized	Parasiti- zation rate percen- tage	Mean No. of parasite per host
<u>Genarchopsis singularis</u> Srivastava, 1933	1	0.277	2.00
<u>Orientocreadium indicum</u> Pande, 1934	3	0.833	2.00
<u>Allocreadium handiai</u> Pande, 1937	2	0.55	1.5
Metacercaria			
<u>Isoparorchis hypselobagri</u> (Billet, 1898) Odhner, 1911	1	0.277	2.00

Channa marulius (Ham.)TABLE 3-C

Parasite	No. of indivi- duals parasi- tized	Parasiti- sation rate percen- tage	Mean No. of parasite per host
<u>Orientocreadum indicum</u> Pande, 1934	1	0.277	3.00
<u>Metaceracaria</u>			
<u>Isoparorchis hypselobagri</u> (Billet, 1898) Odhner, 1911	1	0.277	2.00

Heteropneustes fossilis (B1.)TABLE 3-D

Parasite	No.of indivi- duals parasi- tized	Parasi- tization rate percen- tage	Mean No. of parasite per host
<u>Orientocreadium indicum</u> Pande, 1934	3	0.833	2.33
<u>Orientocreadium pseudobagri</u> Yamaguti, 1934	2	0.55	2.00
<u>Orientocreadium keni</u> n.sp	1	0.277	2.00
<u>Allocreadium handiai</u> Pande, 1937	1	0.277	5.00
<u>Haplorchoides seenghali</u> Dayal and Gupta, 1954	1	0.277	1.00
<u>Phyllodistomum tripathi</u> Motwani and Srivastava 1961	5	1.38	2.4

Clarias batrachus (Linn.)TABLE-3 E

Parasite	No.of indivi- duals parasi- tized	Parasiti- zation rate per- centage	Mean No. of parasite per host
<u>Orientocreadium indicum</u> Pande, 1934	21	5.83	6.047
<u>Orientocreadium pseudobagri</u> Yamaguti, 1934	8	2.22	2.125
<u>Orientocreadium keni</u> n.sp	7	1.94	1.857
<u>Orientocreadium batrachoides</u> Tubangui, 1931	1	0.277	2.00
<u>Allocreadium handiai</u> Pande, 1937	8	2.22	1.875
<u>Pycnadena bariliusi</u> V. Kumari, 1973	1	0.277	1.00

It is evident from the Tables 3A to 3E that O. indicum is the most predominant trematode parasite inhabiting all five host fishes; A. handiai - Parasitized four host sp. viz. C. punctatus, C. striatus, H. fossilis and C. batrachus; O. pseudobagri inhabiting three host species viz. C. punctatus, H. fossilis and C. batrachus; O. keni n.sp. parasitized two host species H. fossilis and Claria batrachus remaining parasites inhabited single host species.

Metacercaria of I. hypsolobagri inhabited all the three species of Channa available locally, whereas other metacercaria lived in single host fish viz. Channa punctatus.

In this table, rate of parasitization and mean number of parasites per host has also been analysed.

Although the results analysed in Tables 3A, 3B, 3C, 3D and 3E give an overall picture of the incidence of trematode infestation found, they do not give an accurate picture of the level of parasitization at any given time of the year. Therefore in tables 4A, 4B, 4C, 4D and in 4E the levels of parasitization of each month of the year are given separately for the five species of host fishes.

LEVEL AND INTENSITY OF PARASITIZATION

LEVEL AND INTENSITY OF PARASITIZATION

Channa punctatus (B1.)

TABLE 4-A

Parasite	July	Aug.	Sept.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	April	May	June
(a) Mean % of fishes parasitized (Level of parasitization)												
<u>Genarchopsis singularis</u> <u>Srivastava, 1933</u>	3.33	3.33	-	-	-	3.33	-	-	-	3.33	-	-
<u>Genarchopsis goppo</u> <u>Srivastava, 1933</u>	-	-	-	-	-	-	-	3.33	-	-	-	-
<u>Genarchopsis piscicola</u> <u>Srivastava, 1933</u>	-	-	3.33	-	3.33	3.33	-	-	3.33	-	-	-
<u>Orientocreadium indicum</u> <u>Pande, 1934</u>	3.33	3.33	-	-	3.33	-	-	-	-	-	-	-
<u>Orientocreadium pseudobagri</u> <u>Yamaguti, 1934</u>	-	6.66	-	-	3.33	-	-	-	-	-	-	-
<u>Allocreadium handiai</u> <u>Pande, 1937</u>	3.33	-	-	-	3.33	3.33	-	-	-	3.33	-	-
Metacercaria												
<u>Euclinostomum heterostomum</u> <u>(Rudolphi, 1809)</u> Travessos, 1928	6.66	-	-	-	-	-	3.33	-	-	-	-	-
<u>Isoparorchis hypselobagri</u> <u>(Billet, 1898)</u> Odhner, 1911	-	-	-	-	-	-	-	-	-	3.33	-	-
<u>Clinostomum complanatum</u> <u>(Rudolphi, 1819)</u> Braun, 1899	10.0	6.66	3.33	3.33	3.33	6.66	6.66	3.33	3.33	3.33	3.33	3.33

Parasite	July	Aug.	Sep.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May	June
(b) Mean number of parasites per host (Intensity of parasitization)												
<u>G. singularis</u> Srivastava, 1933	2.0	2.0	-	-	-	3.0	-	-	-	1.0	-	-
<u>G. goppo</u> Srivastava, 1933	-	-	-	-	-	-	-	2.0	-	-	-	-
<u>G. piscicola</u> Srivastava, 1933	-	-	2.0	-	3.0	2.0	-	-	1.0	-	-	-
<u>O. indicum</u> Pande, 1934	2.0	2.0	-	-	-	6.0	-	-	-	-	-	-
<u>O. pseudobagri</u> Yamaguti, 1934	-	1.5	-	-	1.0	-	-	-	-	-	-	-
<u>A. handiai</u> Pande, 1937	2.0	-	-	-	1.0	1.0	-	-	-	1.0	-	-
<u>E. heterostomum</u> (Rudolphi, 1809) Travassos, 1928	1.5	-	-	-	-	-	1.0	-	-	-	-	-
<u>I. hypselobagri</u> (Billet, 1898) Odhner, 1911	-	-	-	-	-	-	-	-	-	3.0	-	-
<u>Clinostomum complanatum</u> (Rudolphi, 1899) Braun, 1899	3.0	2.0	3.0	3.0	2.0	2.5	2.0	2.0	3.0	2.0	1.0	2.0

Channa striatus (B1.)

TABLE 4-B

Parasite	July	Aug.	Sep.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	April	May	June
(a) Mean % of fishes parasitized (Level of parasitization)												
<u>Generchoopsis singularis</u> Srivastava, 1933	-	-	-	3.33	-	-	-	-	-	-	-	-
<u>Orientocreadium indicum</u> Pande, 1934	3.33	-	3.33	-	3.33	-	-	-	-	-	-	-
<u>Allocreadium handiai</u> Pande, 1937	-	3.33	-	-	-	-	-	-	-	3.33	-	-
Metacercaria												
<u>Isoparorchis hypselobagri</u> (Billet, 1898) Odhner, 1911	-	-	-	-	-	-	3.33	-	-	-	-	-

Parasite	July	Aug.	Sep.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	April	May	June
(b) Mean number of parasites per host (Intensity of Parasitization)												
<u>G. singularis</u> Srivastava, 1933	-	-	-	2-0	-	-	-	-	-	-	-	-
<u>O. indicum</u> Pande, 1934	3.0	-	2.0	-	1.0	-	-	-	-	-	-	-
<u>A. handiai</u> Pande, 1937	-	2.0	-	-	-	-	-	-	-	1.0	-	-
<u>I. hypselobagri</u> (Billet, 1898) Odhner, 1911	-	-	-	-	-	-	2.0	-	-	-	-	-

Channa marulium (Ham.)

TABLE 4-C

Parasite	July	Aug.	Sep.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May	June
(a) Mean % of fishes parasitized (Level of parasitization)												
<u>Orientocreadium indicum</u> Pande, 1934	-	3.3	-	-	-	-	-	-	-	-	-	-
Metacercaria												
<u>Isoparorchis hypselobaqui</u> (Billet, 1898, Odhner, 1911)	-	-	3.33	-	-	-	-	-	-	-	-	-

Parasite	July	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar.	Apl.	May	June
(b) Mean number of parasites per fish (Intensity of parasitization)												
<u>O. indicum</u>												
Pande, 1934	-	3.0	-	-	-	-	-	-	-	-	-	-
<u>I. hypselobagri</u>												
(Billet, 1898)	-	-	2.0	-	-	-	-	-	-	-	-	-
Odhner, 1911												

Heteropneustes fossilis (B1.)

TABLE 4-D

Parasite	July	Aug	Sep	Oct	Nov.	Dec	Jan	Feb	Mar	Apr	May	June
(a) Mean % of fishes parasitized (Level of parasitization)												
<u>Orientocreadium indicum</u> Pande, 1934	3.33	-	-	-	3.33	-	-	-	3.33	-	-	-
<u>Orientocreadium pseudobagri</u> Yamaguti, 1934	-	-	-	-	-	-	-	6.66	-	-	-	-
<u>Orientocreadium keni</u> n.sp.	-	-	-	-	-	-	3.33	-	-	-	-	-
<u>Allocreadium handiai</u> Pande, 1937	-	-	-	-	-	-	-	3.33	-	-	-	-
<u>Haplorchoides seenghali</u> Dayal & Gupta, 1954	-	-	-	-	-	-	-	3.33	-	-	-	-
<u>Phyllodistomum tripathi</u> Motwani & Srivastava, 1961	6.66	10.0	-	-	-	-	-	-	-	-	-	-

Parasite	July	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	June
(b) Mean number of parasites/fish (Intensity of parasitization) per												
<u>O. indicum</u> Pande, 1934	2.0	-	-	-	2-0	-	-	-	3.0	-	-	-
<u>O. pseudobagri</u> Yamaguti, 1934	-	-	-	-	-	-	-	2.0	-	-	-	-
<u>O. keni</u> n.sp.	-	-	-	-	-	-	2.0	-	-	-	-	-
<u>A. handiai</u> Pande, 1937	-	-	-	-	-	-	-	5.0	-	-	-	-
<u>H. seenghali</u> Dayal & Gupta, 1954	-	-	-	-	-	-	-	1.0	-	-	-	-
<u>P. tripathi</u> Motwani & Srivastava, 1961	2.5	2.33	-	-	-	-	-	-	-	-	-	-

Clarias batrachus (Linn.)

TABLE 4-E

Parasite	July	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	April	May	June
(a) Mean % of fishes parasitized (Level of parasitization)												
<u>Orientocreadium indicum</u> Pande, 1934	-	6.66	6.66	6.66	3.33	10.0	10.0	6.66	6.66	3.33	-	-
<u>Orientocreadium pseudobagri</u> Yamaguti, 1934	-	3.33	6.66	3.33	3.33	-	3.33	3.33	3.33	-	-	-
<u>Orientocreadium keni</u> n.sp.	-	6.66	-	-	-	-	3.33	6.66	3.33	-	-	-
<u>Orientocreadium batrachoides</u> Tubangui, 1931	-	-	-	-	-	-	-	-	-	3.33	-	-
<u>Allocreadium handiai</u> Pande, 1937	-	-	-	3.33	3.33	-	-	6.66	6.66	6.66	-	-
<u>Pycnadena barilliusi</u> Kumari, 1973	-	-	-	-	-	-	3.33	-	-	-	-	-

Parasite	July	Aug	Sep.	Oct	Nov	Dec	Jan	Feb	Mar	April	May	June
(b) Mean number of parasites per host (Intensity of parasitization)												
<u>O. indicum</u> Pande, 1934	-	8.0	7.5	7.5	8.0	4.33	8.33	3.0	1.5	2.0	-	-
<u>O. pseudobagri</u> Yamaguti, 1934	-	3.0	2.5	3.0	2.0	-	2.0	1.0	1.0	-	-	-
<u>O. keni</u> n.sp.	-	2.0	-	2.0	-	-	1.0	1.0	4.0	-	-	-
<u>O. batrachoides</u> Tubangu, 1931	-	-	-	-	-	-	-	-	-	2.0	-	-
<u>A. handiai</u> Pande, 1937	-	-	-	1.0	1.0	-	-	2.5	2.0	2.0	-	-
<u>P. bariliusi</u> Kumari, 1973	-	-	-	-	-	-	1.0	-	-	-	-	-

The analysis of the above Tables (4A, 4B, 4C, 4D & 4E) shows that in Channa punctatus (Table 4A) the highest incidence of parasitization by G.singularis is found uniformly in the month of July, August, December and April; by G.piscicola the highest incidence of parasitization in September, November, December and March; Allocreadium handiai had the highest incidence of parasitization in July, November, December & April and Clinostomum complanatum metacercaria were found uniformly in all months with highest incidence of parasitization in the month of July and minimum level of parasitization during the months of September, October, November, February, March, April, May and June.

In Channa striatus (Table 4-B) the highest incidence of parasitization is found uniformly by O.indicum in the month of July, September and November.

In Channa marulius (Table 4-C) the incidence of parasitization by O.indicum is found in the month of August, by Isoparorchis hypselobagri (metacercaria) the incidence is found in the month of September.

In H.fossilis (Table 4-D) the highest incidence of parasitization by P.tripathi was found in the month of August and lowest in July.

In Clarias batrachus (Table 4-E) the highest incidence of parasitization by O.indicum was found in the month of August and lowest in November and April.

SEASONAL INCIDENCE OF PARASITES

SEASONAL INCIDENCE OF PARASITES

Channa punctatus (B1.)

TABLE 5-A

Parasites	Winter season (N D J F)		Summer season (M A M J)		Rainy season (J A S O)	
	(a)	(b)	(a)	(b)	(a)	(b)
<u>Genarchopsis singularis</u> Srivastava, 1933	0.8325	0.750	0.8325	0.25	1.665	1.00
<u>Genarchopsis goppo</u> Srivastava, 1933	0.8325	0.5	-	-	-	-
<u>Genarchopsis piscicola</u> Srivastava, 1933	1.665	1.25	0.8325	0.25	0.8325	0.50
<u>Orientocreadium indicum</u> Pande, 1934	0.8325	1.5	-	-	1.665	1.00
<u>Orientocreadium pseudobagri</u> Yamaguti, 1934	0.8325	0.25	-	-	1.665	0.375
<u>Allocreadium handiai</u> Pande, 1937	1.665	0.5	0.8325	0.25	0.8325	0.50
Metacercaria						
<u>Euclinostomum heterostomum</u> (Rudholphi, 1809), Travassos, 1928	0.8325	0.25	-	-	1.665	0.375
<u>Isoparorchis hypselobagri</u> (Billet, 1898) Odhner, 1911	-	-	0.8325	0.75	-	-
<u>Clinostomum complanatum</u> (Rudolphi, 1819) Braun, 1899	4.995	2.125	3.3300	2.0	5.830	2.75

Channa striatus (B1.)

TABLE 5-B

Parasites	Winter season		Summer season		Rainy season	
	(N D J F) (a)	(b)	(M A M J) (a)	(b)	(J A S O) (a)	(b)
<u>Genarchopsis singularis</u> Srivastava, 1933	-	-	-	-	0.8325	0.5
<u>Orientocreadium indicum</u> pande, 1934	0.8325	0.25	-	-	0.8325	1.25
<u>Allocreadium handiai</u> pande, 1937	-	-	0.8325	0.25	0.8325	0.5
Metacercaria						
<u>Isoparorchis hypselobagri</u> (Billet, 1898) Odhner, 1911	0.8325	0.5	-	-	-	-

(a) % of hosts parasitized

(b) Mean No. of parasites per host

Channa marulius (Ham.)TABLE 5-C

Parasites	Winter season (N D J F)		Summer season (M A M J)		Rainy season (J A S O)	
	(a)	(b)	(a)	(b)	(a)	(b)
<u>Orientocreadium indicum</u> Pande, 1934	-	-	-	-	0.8325	0.75
Metacercaria						
<u>Isoparorchis hypselobagri</u> (Billet, 1898) Odhner, 1911	-	-	-	-	0.8325	0.5

(a) % of hosts parasitized

(b) Mean No. of parasites per host.

Heteropneustes fossilis (B1.)

TABLE 5-D

Parasites	Winter season (N D J F)		Summer season (M A M J)		Rainy season (J A S O)	
	(a)	(b)	(a)	(b)	(a)	(b)
<u>Orientocreadium indicum</u> Pande, 1934	0.8325	0.5	0.8325	0.75	0.8325	0.5
<u>Orientocreadium pseudobagri</u> Yamaguti, 1934	1.665	0.5	-	-	-	-
<u>Orientocreadium keni</u> n.sp.	0.8325	0.5	-	-	-	-
<u>Allocreadium handiai</u> Pande, 1937	0.8325	1.25	-	-	-	-
<u>Haplorchoides seenghali</u> Dayal and Gupta, 1954	0.8325	0.25	-	-	-	-
<u>Phyllodistomum tripathi</u> Motwani and Srivastava, 1961	-	-	-	-	4.165	1.275

(a) % of hosts parasitized

(b) Mean No. of parasites per host.

Clarias batrachus (Linn.)

TABLE 5-E

Parasites	Winter season (N D J F)		Summer season (M A M J)		Rainy season (J A S O)	
	(a)	(b)	(a)	(b)	(a)	(b)
<u>Orientocreadium indicum</u> Pande, 1934	7.4975	5.915	2.4975	0.875	7.495	5.75
<u>Orientocreadium pseudobagri</u> Yamaguti, 1934	2.4975	1.25	0.8325	0.25	3.33	2.125
<u>Orientocreadium keni</u> n.sp.	2.4975	0.5	0.8325	1.0	2.4975	1.0
<u>Orientocreadium batrachoides</u> Tubangui, 1931	-	-	0.8325	0.5	-	-
<u>Allocreadium handiai</u> Pande, 1937	2.4975	0.875	3.33	1.0	0.8325	0.25
<u>Pycnadena bariliusi</u> V. Kumari, 1973	0.8325	0.25	-	-	-	-

(a) % of hosts parasitized

(b) Mean No. of parasites per host.

It is clear from the Table 5-A that in Channa punctatus the maximum infestation is recorded during winter season. The highest percentage of the fish, Channa punctatus parasitized by G. singularis and the highest mean number of parasites per host are in rainy season. The highest percentage of hosts parasitized by G. piscicola and the highest mean number of parasites per host occurred during winter season. The highest percentage of hosts parasitized by A. handiai occurred in winter and highest mean number of parasites per host in rainy and winter season. The highest percentage of the fish, C. punctatus parasitized by Clinostomum complanatum (metacercaria) and the highest mean number of parasites per host occurred during rainy season.

In Channa striatus (Table 5-B) the maximum number of parasites were recorded in rainy season. The highest percentage of hosts parasitized by O. indicum occurred during winter and rainy season and the highest mean number of parasites per host during rainy season. The highest percentage of host parasitized by A. handiai occurred in summer and rainy season and highest mean number of parasites per host in rainy season.

In Channa marulius (Table 5-C) the maximum number of parasites were recorded in rainy season. The high percentage of the fish Channa marulius parasitized by O. indicum and the highest mean number of parasites per host are recorded in

rainy season. The highest percentage of the fish, Channa marulius parasitized was by Isoparorchis hypselobagri (metacercaria) and the highest mean number of parasites per host are recorded in rainy season.

In H.fossilis (Table 5-D) the maximum number of parasites were recorded during winter season. The high percentage of hosts parasitized by O.indicum occurred during winter, summer and rainy season and the highest mean number of parasites per host during summer season.

In Clarias batrachus (Table 5-E) the maximum number of trematodes have been recorded during the winter season. The highest percentage^t_x of fishes parasitized by O.indicum and the highest mean number of parasites per host occurred during winter season. The highest percentage of this fish host parasitized by O.pseudobagri and the highest mean number of parasites per host occurred during rainy season. The highest percentage of fishes parasitized by O. keni n.sp. occurred during winter and rainy season and highest mean number of parasites per host in summer and rainy season. The highest percentage of fishes parasitized by A.handiai and the highest mean number of parasites per host occurred during summer season.

It is clear from the study of above tables (Table 5-A, 5-B, 5-C, 5-D, and 5-E) that seasonal incidence of parasites was higher in all host fishes during rainy season. Out of five host fishes, C. batrachus, C. punctatus, C. striatus and H. fossilis have infection in all seasons, while the least infected host fish C. marulius (Ham.) was found infected only during rainy season. Thus seasonal prevalence forms one of the major ecological factor influencing the incidence of helminth infection in host fishes.

- - - - -

EXPLANATION OF LIST AND TABLES

	<u>Page</u>
List O-1 Host-Parasite List	16 - 19
List O-2 Parasite - Host List	20 - 22
Table 1 Showing the index of Total Helminth Infection (Trematodes)	74
Table 2-A Shows the Hostwise Analysis of Parasites in <u>Channa punctatus</u> (Bl.)	76 - 77
Table 2-B Shows the Hostwise Analysis of Parasites in <u>Channa striatus</u> (Bl.)	78
Table 2-C Shows the Hostwise Analysis of Parasites in <u>Channa marulius</u> (Ham.)	79
Table 2-D Shows the Host-wise Analysis of Parasites in <u>Heteropneustes</u> <u>fossilis</u> (Bl.)	80
Table 2-E Shows the Host-wise Analysis of Parasites in <u>Clarias batrachus</u> (Linn.)	81
Table 3-A Shows the overall Incidence of Trematodes in <u>Channa punctatus</u> (Bl.)	84
Table 3-B Shows the overall Incidence of Trematodes in <u>Channa striatus</u> (Bl.)	85
Table 3-C Shows the overall Incidence of Trematodes in <u>Channa marulius</u> (Ham.)	86
Table 3-D Shows the overall Incidence of Trematodes in <u>Heteropneustes</u> <u>fossilis</u> (Bl.)	87
Table 3-E Shows the overall Incidence of Trematodes in <u>Clarias batrachus</u> (Linn.)	88
Table 4-A Shows the level and Intensity of Parasitization in <u>Channa punctatus</u> (Bl)	90-91
Table 4-B Shows the level and Intensity of Parasitization in <u>Channa striatus</u> (Bl)	92-93

Table 4-C	Shows the level and intensity of Parasitization in <u>Channa marulius</u> (Ham.)	94 - 95
Table 4-D	Shows the level and intensity of Parasitization in <u>Heteropneustes fossilis</u> (Bl.)	96 - 97
Table 4-E	Shows the level and intensity of Parasitization in <u>Clarias batrachus</u> (Linn.)	98 - 99
Table 5-A	Shows the seasonal incidence of parasites in <u>Channa punctatus</u> (Bl.)	101
Table 5-B	Shows the seasonal incidence of parasites in <u>Channa striatus</u> (Bl.)	102
Table 5-C	Shows the seasonal incidence of parasites in <u>Channa marulius</u> (Hamm.)	103
Table 5-D	Shows the seasonal incidence of parasites in <u>Heteropneustes fossilis</u> (Bl.)	104
Table 5-E	Shows the seasonal incidence of parasites in <u>Clarias batrachus</u> (Linn.)	105

- - - - -

EXPLANATION OF PLATES AND FIGURESPLATE 1

Fig. 1 Pycnadena bariliusi Kumari, 1973 (Ventral view)

PLATE 2

Fig. 2 Allocreadium handiai Pande, 1937 (Ventral view)

PLATE 3

Fig. 3 Orientocreadium batrachoides Tubangui, 1931
(Ventral view)

PLATE 4

Fig. 4 Orientocreadium indicum Pande, 1934
(Ventral View)

PLATE 5

Fig. 5 Orientocreadium pseudobagri Yamaguti, 1934
(Ventral View)

PLATE 6

Fig. 6 Orientocreadium keni n.sp. (Ventral View)

PLATE 7

Fig. 7 Phyllodistomum tripathi Motwani & Srivastava, 1961
(Ventral View)

PLATE 8

Fig. 8 Metacercaria of Isoparorchis hypselobagri
(Billet, 1898) Odhner, 1911
(Ventral View)

PLATE 9

- Fig. 9 Genarchopsis piscicola Srivastava, 1933
(Ventral View)

PLATE 10

- Fig. 10 Genarchopsis goppo Srivastava, 1933
(Ventral view)

PLATE 11

- Fig. 11 Genarchopsis singularis Srivastava, 1933
(Ventral view)

PLATE 12

- Fig. 12 Metacercaria of Euclinostomum heterostomum
(Rudolphi, 1809) Travassos, 1928
(Ventral view)

PLATE 13

- Fig. 13 Haplorchoides seenghali Dayal and Gupta, 1954
(Ventral view)

PLATE 14

- Fig. 14 Graph showing the helminth infection (Trematodes)
in the fishes.

PLATE 15

- Fig. 15 Histogram showing the overall incidence of
Trematodes in the fish, Channa punctatus (Bl.)

PLATE 16

- Fig. 16 Histogram showing the overall incidence of
Trematodes in the fish, Channa striatus (Bl.)

PLATE 17

Fig. 17 Histogram showing the overall incidence of Trematodes in the fish, Channa marulius (Ham.)

PLATE 18

Fig. 18 Histogram showing the incidence of five species of the Trematodes in the fishes, Heteropneustes fossilis (Bl.)

PLATE 19

Fig. 19 Histogram showing the overall incidence of six species of the Trematodes in the fish, Clarias batrachus (Linn.)

- - - - -




Fig. 1.

Ryncadena bariliusi . Kumari, 1973; Ventral view.

PLATE 2.




Fig. 2.

Allocreadium handiai Pande, 1937; Ventral view.

PLATE 3.




Fig. 3.

Orientocreadium batrachoides Tubangui, 1931; Ventral view

PLATE 4.




Fig. 4.

Orientocreadium indicum pande, 1934; Ventral view




Fig. 5

Orientocreadium pseudobagri Yamaguti, 1934; Ventral view.

PLATE 6.




Fig. 6

Orientocreadium keni n.sp. Ventral view.

PLATE 7.




Fig. 7.

Phyllodistomum tripathi Motvani and Srivastava,
1961; Ventral view.

PLATE 8.




Fig. 8.

Metacercaria of Isoparorchis hypselobagri
(Billet, 1898) Odhner, 1911; Ventral view.

PLATE 9.




Fig. 9.

Genarchopsis piscicola Srivastava, 1933; Ventral view.




Fig. 10 ,

Genarchopsis goppo Srivastava, 1933; Ventral view.

PLATE 11




Fig. 11

Genarchopsis singularis Srivastava, 1933; Ventral view




Fig. 12.

Metacercaria of Euclinostomum heterostomum

(Rudolphi, 1809) Travassos, 1928; Ventral view.

PLATE 13.




Fig. 13

Haplorchooides seenghali Dayal and Gupta, 1954

Ventral view.

PLATE 14.

Index of Helminth infection (trematodes)
July 1993 to June 1995





Fig. 14

PLATE 15.

Over all Incidence of Trematodes

July 1993 to June 1995




In *Channa punctatus* (Bl.)

Fig. 15

PLATE 16.

Over all Incidence of Trematodes

July 1993 to June 1995




In Channa striatus (Bl.)

Fig. 16.

PLATE 17.

Over all Incidence of Trematodes

July 1993 to June 1995




In Channa marulius (Ham.)

Fig. 17.

PLATE 18.

Over all Incidence of Trematodes

July 1993 to June 1995




In *Heteropneustes fossilis* (Bl.)

Fig. 18.

PLATE 19.

Over all Incidence of Trematodes

July 1993 to June 1995



In Clarias batrachus (Linn.)

Fig. 19.

S U M M A R Y

SUMMARY

To investigate helminth fauna from some fresh water fishes of Bhundelkhand region, the author under the present project has planned to examine five host species of fresh water air-breathing fishes, most commonly available in this region. These fishes include Clarias batrachus (Linn.); H. fossilis (B1.) and Channa sp. namely Channa punctatus (B1.) Channa striatus (B1.) and Channa marulius (Ham.). Out of five species of Channa available from Uttar Pradesh viz - Channa gachua (Ham.); Channa leucopunctatus (B1.); Channa striatus (B1.); Channa punctatus (B1.) and Channa marulius (Ham.) only above mentioned three species were available from water bodies of Bundelkhand region, including Jhansi, hence they have been included in the present study. These fishes were collected regularly from Upper Dam, Lower Dam, Chuna bhatti tank, Kolas pond and other fresh water bodies in and around Jhansi and from local fish markets for a period of two years from July 1993 to June 1995. On an average fifteen fishes of each species (Total 75 fishes per month) were collected, examined and dssected. A total of 1800 fishes were examined during a period of two years. A thorough search was done to collect the parasites from various organs of fishes.

Soon after collection the trematodes were studied alive to observe spines or papillae if any, excretory system and genital opening. They were then fixed in 5-10% formaline, after fixation and thorough washing in water, worms were dehydrated in alcoholic series, then stained in borax carmine, cleared in xylene and finally mounted in DPX.

For preparing whole mount of trematodes, the worms were relaxed in fresh water for some time. The body was gently stretched by adding luke warm water. Specimens were kept in 5 - 10% formaline overnight, then washed with water, kept in alcoholic series, stained in acetoalum carmine, cleared in clove oil and finally mounted in canada balsum/dpx.

The work incorporated in the present thesis is divided into three parts.

Part I deals with introduction, historical resume, material and method, a systematic list of host examined, host-parasite list and parasite-host list.

Part II deals with the taxonomic grouping of thirteen trematodes including three metacercarial forms recovered during the study period. Mention has not been made of nematodes or cestodes as these were

not the part of project. Out of these trematode species, one species has been described as new species where as remaining ten are redescribed in detail, furnishing further information and observations which were essential to enrich our knowledge on these parasites. They are known forms but majority of these form the first host and locality record. These trematodes belong to eight families of the order Digenea and include :

1. Pycnadena bariliusi Kumari, 1973 has been described from the stomach of Clarias batrachus (Linn.). It is characterised by long hind body, symmetrical testes, posttesticular extension of uterus and embryonated eggs with oculate miracidia and absence of postoral ring.

It forms the first host and locality record.

2. Allocreadium handiai Pande, 1937 has been recorded from the intestine of four host species excluding Channa marulius (Ham.). It is characterised by the shape of the body, acetabulum smaller than oral sucker, tandem testes and small ovary attached to acetabulum.

It is the first host and locality record from Bhundelkhand region.

3. Orientocreadium batrachoides Tubangui, 1931 has been collected from the intestine of Clarias batrachus (Linn.). It is characterised by receptaculum seminis, pretesticular, equatorial ovary and postequatorial testes. Vesicula seminalis externa is long and saccular.

This recorded form has been described from this region.

4. Orientocreadium indicum Pande, 1934 has been described from the intestine of all the five host species under the present project. It is characterised by the presence of spined cirrus and metraterm, spherical ovary, vitellaria extending from posterior margin of acetabulum to posterior end of body and the follicles of two sides meet without forming lattice.

It forms the first host and locality record.

5. Orientocreadium pseudobagri Yamaguti, 1934 has been recorded from the intestine of two host species viz - Clarias batrachus (Linn.) and Heteropneustes fossilis (Bl.). It is characterised by oval oral sucker, small prepharynx, long oesophagus and extension of vitellaria halfway between the caudal testis and posterior tip of the body and cirrus pouch on the right side of acetabulum instead of overlapping it.

It forms the first host and locality record.

6. Orientocreadium keni n.sp. has been collected from the intestine of two host species viz Clarias batrachus (Linn.) and Heteropneustes fossilis (Bl.). This new species is characterised by small body, transversely elongated ovary and testes, very small oesophagus and eggs rounded or oval.

It forms the first host and locality record.

7. Phyllodistomum tripathi Motwani and Srivastava, 1961 has been recorded from the intestine of Heteropneustes fossilis (Bl.). It is characterised by acetabulum distinctly larger than oral sucker, wavy margins having thick folds along with margins, flask shaped body, divisible into a narrow, tubular fore body and a foliate hind body. Prepharynx and pharynx absent, deeply lobed testes, cirrus sac absent, lobed ovary, overlaps the right vitellaria. Receptaculum seminis absent, vitellaria two lobed glands.

It forms the first host and locality record.

8. Isoparorchis hypselobagri (Billet, 1898) Odhner 1911 has been recorded from the body cavity of all the three host species of Channa (Bl.). It is characterised by body with thick cuticle, intestinal caeca long and serpentine. Acetabulum large than oral sucker.

This recorded form has been described from this region.

9. Genarchopsis piscicola Srivastava, 1933 has been recorded from the intestine of Channa punctatus (B1.). It characterised by small and fusiform body. Oral sucker cuplike, acetabulum large and well developed, prepharynx and oesophagus absent. Cirrus sac absent, uterine coils intercaecal, extending posteriorly upto the hind border of the testes.

This recorded form has been described from Bhundelkhand region.

10. Genarchopsis goppo Srivastava, 1933 has been collected from the intestine of Channa punctatus (B1.). It is characterised by elliptical shape of the body, acetabulum large and spherical. Cirrus sac absent. Vesicula seminalis well developed, vitellaria two lobed glands.

It forms the first host and locality record.

11. Genarchopsis singularis Srivastava, 1933 has been described from the intestine of Channa punctatus (B1.) and Channa striatus (B1.). It is characterised by small oesophagus, symmetrical testes, cirrus sac absent and two compact vitellaria.

The parasite forms the first host and locality record.

12. Euclinostomum heterostomum (Rudolphi, 1809)

Travassos, 1928 has been collected from the body cavity of Channa punctatus (B1.). It is characterised by presence of well developed pharynx and vitellaria, body large with collar like formation at anterior end. Intestinal caeca very thin upto posterior border of acetabulum with lateral diverticulae. Vitelline follicles very small and immature.

This recorded form has been described from this region.

13. Clinostomum complanatum (Rudolphi, 1819) Braun, 1899

simple record of this metacercaria is mentioned. This metacercaria was recovered in large numbers throughout the study period from July 1993 to June 1995 from the body cavity of Channa punctatus (B1.).

14. Haplorchoides seenghali Dayal and Gupta, 1954 has been recorded from the intestine of H. fossilis (B1.). It is characterised by small size of body, large size of testis, absence of cirrus pouch and acetabulum.

It forms the first host and locality record.

Part III deals with host parasite relationship. An attempt has been made to correlate various ecological factors-biotic and abiotic, affecting host-parasite relationship. The data collected for a period of two years from

July 1993 to June 1995 has been statistically analysed to study index of total helminth infection, host-wise analysis, overall incidence, level and intensity of parasitization, seasonal incidence, incidence of infection in relation to sex, co-relation between the size and sex of fish and intensity of infection in trematodes.

In the end of thesis, the references of the literature cited are given. Further, a brief 'Summary' of the work done and included in the thesis is also added in the end.

The present thesis extends over 173 pages and is illustrated with thirteen plates containing camera lucida diagrams of the trematode species described. Beside this one graph and five histograms are also given. All these figures are original and drawn by the author. The thesis includes two 'Lists' and 20 Tables which presents the datas collected during the entire work.

- - - - -

REFERENCES

REFERENCES

- AGRAWAL, P.K. AND AGRAWAL, S.M., 1981 - Life history of Haplorchoides vacha n.sp. (Trematode : Heterophyidea) from the intestine of Eutropiichthys vacha (Ham.) Riv. Parasite XLII (1).
- AGRAWAL, S.M., 1959 - Further observations on Euclinostomum indicum Bhalerao, 1942 (Trematoda : Clinostomidae). Ind. J. Helm. 10, 31-39.
- AGRAWAL, V. 1964 - Studies on some new trematodes from fresh water fishes of Lucknow. Ind. J. Helm. 16(2) 82-89.
- AGRAWAL, V., 1964 - On some new trematodes from fresh water fishes of Lucknow. Indian J. Helminth. 31, 219-222.
- AGRAWAL, V., 1966 - Studies on some trematode parasites of fresh water fishes from Lucknow. Ann. Parasit 41 (3), 217-231.
- AHMAD J. 1978 - Digenetic trematodes from marine fishes. Geobios, 5, 281-283.
- AHSAN, S.M., 1959 - On the occurrence of metacercaria of Clinostomum sp. in Trichogaster fasciatus. Curr. Sci. 1, 32-33.

ANANTARAMAN, M. 1959 - Metacercaria (Fillocreadoides) in the planktonic ctenophore, Pleurobranchia globosa Moser, 1903, from the Madras Coast. Nature, 183, 1407-1408.

ANJANEYULU, G., 1967 - Observations on the femal reproductive system of Genarchopsis punctati Agrawal, 1966, Ind. J. Helminth., 19(2), 118-121.

BASHIRULLAH, A.K.M. & MUSTAK ELAHI., 1972 - On two new species of Genarchopsis Ozaki, 1925 from fresh water fishes of Dacca, Bangladesh. Riv. Parasit., XXXIII (4), 277-280.

BASHIRULLAH, A.K.M. & HAFIZUDDIN, A.K.M., 1976 - Digenetic trematodes from fresh water fishes of Bangladesh. Riv. Parasit., XXXVII (1), 35-39.

BAUGH, S.C. 1956 - Contribution to our knowledge of digenetic trematodes III. Proc. Nat. Acad. Sci. India, 26, 295-301.

BAUGH, S.C., 1958 - Contribution to our knowledge of digenetic trematodes III. Proc. Nat. Acad. Sci. India, 28, 205-226.

BAUGH, S.C., 1962 - Contribution to our knowledge of Digenetic trematodes. VI. Zeit. Fur. Parasitenk., 22, 303 - 315.

- BAUGH, S.C. and PANDEY, K.C., 1969 - Studies on Clinostome metacercariae, A restudy of Clinostomum giganaticum. Ang. Parasitol., 10, 211-233.
- BEAR, K.E. Von, 1827 - Beitrage zur kenntnis der neideren Thiere Nova Acta Leop. Carol, 13, 523-762.
- BEVERLEY-BURTON, M. 1962 - Some trematodes from Clarias spp. in the Phodesias, including Allocreadium mazoensis n.sp. and Eumaseia bangwelensis n.sp. and comments on the species of the genus Orientocreadium Tubangui, 1931. Proc. Helm. Soc. Wash., 29, 103-115.
- BHADAURIA, S. & DANDOTIA, M.R. 1984 - Studies on the digenetic trematodes of fresh water fishes with special reference to Gwalior region, Part III. Seasonal incidence of some Treematode Parasites of food fishes of Gwalior region. Rivista di Parassitologia, Vol. I (XLV), 419-429.
- BHADAURIA, S. & DANDOTIA, M.R. 1992 - Studies on the Trematode Parasites of fresh water fishes with special reference to Gwalior region. Part II. On one new Genus and some unknown and known species. Rivista di Parassitologia, Vol. IX 341 - 383.

BHADAURIA, S. & DANDOTIA, M.R. 1994 - Studies on the digenetic Trematodes of fresh water fishes with special reference to Gwalior region. Part IV. Description of ten new and six already known forms belonging to eight Genera of trematodes. Rivista di Parassitologia, Vol. XI. (XLVII), 353 - 397.

BHALERAO, G.D. 1926 - On two synonym of the genera Isoparorchis Southwell, 1913 and Leptolecithum Kobayashi 1920, with a description of the male genitalia of Isoparorchis trisimilitubis Southwell, 1913. Ann. Mag. Nat. Hist., 17, 246-250.

BHALERAO, G.D. 1932 - A note on the Probability of infection of man and domestic carnivores by Isoparorchis hypsalobagri (Billet, 1898). Indian J. Vet. Sci., 2, 406-407.

BHALERAO, G.D. 1936 - Studies on helminths of India, Trematodes II. J. Helminth., 14, 181-206.

BHALERAO, G.D. 1936 - Studies on the helminths of India. Trematoda III. J. Helminth., 14, 207-228.

BHALERAO, G.D. 1937, - Studies on the helminths of India. Trematoda IV. J. Helminth., 15(2), 97-124.

BHALERAO, G.D. 1942 - Some metacercarial forms of Clinostom-
tides (Trematoda) from India. Proc. Indian
Acad. Sci., (Sect. B) 16, 67-71.

BHALERAO, G.D. 1942 - Rec.Ind. Mus. 44, 191-192.

BHALERAO, G.D. 1943 - On two trematodes from fishes in India.
Proc. Indian Acad. Sci., 18, 119-124.

BHATNAGAR, G.P., 1986 - Limnology of lower lake of Bhopal (M.P)

BOVBIEN, P. 1926 - Caryophyllaeidae from Java. Videnskabelige
Meddeleser fra Dansk naturhistorisk forening
L. Kobehnavn, 82, 157-181.

BRAUN, M. 1899 - Uber Clinostomum Leidy. Zool.Anz., 22, 489-493.

BRAUN, M. 1900 - Die Arten der Gattung Clinostomum Leidy.
Zool. Jahrb. Syst., 14 (1), 1-48.

BYRD, E.E., UERNARD, G.E. AND REIBER, R.S., 1940 - The excretory
system in trematode. I. Studies on the
excretory system in trematode sub-family
Gorgoderinae looss, 1899 J. Parasit., 26, 407-420.

CANNING, E.U. AND MADHVI, R. 1977 - Studies on two new species
of Microsporida hyperparasite in adult
Allocreadium fasciatusi (Trematoda, Allocrea-
dium), Parasitol., 75, 293-300.

- CHANDLER, A.C. 1916 - A study of the structure of feathers,
with reference to their taxonomic significance.
Univ. Calif. Pub. Zool., 13 (11), 243-446.
- CHATTERJI, R.C. 1933 - On the trematode parasites of a Rangoon
siluroid fish Clarias batrachus (Linn., 1785).
Bull. Acad. Sci. Allahabad 3(1), 33-40.
- CHATTERJI, R.C. 1948 - On some heterophyid trematodes of the
genus Haplorchis Looss 1899. Abstract. Proc.
34th Ind. Sc. Cong., 175.
- CHATTERJI, P.N. 1953 - Two new Heterophyid trematodes of the
genus Haplorchis Looss 1899, Nat. Acad. Sci.
India, 23 (IV-VI), 163-174.
- CHATTERJI, P.N. 1956 - Two new fish trematodes of the family
Heterophyidae Odhner, 1914. Proc. Nat. Acad.
Sci. India, (B) 26(5), 247-355.
- CHAUHAN, B.S. 1945 - Trematodes from Indian marine fishes.
Part IV. on some trematodes of the family
Hemiuridae Luhe, 1901 with description of six
new forms. Proc. Indian Acad. Sci. 21, 160-173.
- CHAUHAN B.S. 1947 - Notes on some Helminths in the collection
of the Zoological Survey of India. Rec.
Indian Musc., 45, 133-137.

- CHAUHAN, B.S. 1949 - Fish and fisheries of the Patna State, Orissa, Rec. Indian Mus., 45, 267-282.
- CHAUHAN, B.S. 1953 - Studies on the trematode fauna of India IV. Sub-class Digenea (Prostomata) A reivision of Hemiuroidea from the Indian region. Rec. Indian Mus. , 51, 289-293.
- CHAUHAN, B.S. 1955 - On the taxonomic position and distribution of the trematode genus Isoparorchis Southwell, 1913. J. Zool. Soc. India, 7, 87-90.
- CHAUHAN, B.S. 1954 - Studies on the trematode fauna of India. Part II, Sub class Aspidogastrea, Rec. Ind. Mus. , 51, 209-230.
- CHAUHAN, B.S. 1957 - Two new fish trematodes of the family Heterophyidae Odhner, 1914. Proc. Nat. Acad. India, 26, 347-355.
- CHAUHAN, B.S., 1963 - Helminthological Studies in India. Proc. 50th Indian Sci. Congr., Part-II, Presidential address, 174-185.
- CHAKRABORTI, K.K., 1970 - A double metacercarial infection in an Indian fresh water fish, Heteropneustes fossilis (Bloch). Rev. Biol. Trop. Med. Parasit., 17, 91-96.

- CHAKRAVARTY, R. & TANDON, V., 1988 - On the present status of caryophyllidea with a fresh water catfish Clarias batrachus (L.) in north-east India and a record of an Anomalous form. Indian J. Helminth. (N.S.), 5 (1), pp. 37-54.
- CHEN, H.T., 1936 - A study of the Haplorchinae (Looss, 1899) Poche, 1926 (Trematoda : Heterophyidae). Parasit., 28, 40-55.
- CHEN, H.T., 1949 - Systematic consideration of some heterophyid trematodes in the subfamilies Haplorchinae and Stellantchasminae. Annals Trop. Med. Parasit., 43, 304-312.
- CHINDERMAN, C.J., 1953 - Fisheries report for lakes of North Central Massachusetts (Div. Fish & Game, Dept. Conservation) Boston, (1950), 1-28.
- COHN, L., 1903 - Helminthologische Mitteilungen. Arch. Naturgesch., 69, 47-68.
- COI, W.H. & KUNTZ, R.E. 1960 - Trematode parasites of vertebrates of East Pakistan. Trans. Am. Microsc. Soc., 79 (3), 281-286.

DANNIS, E.A. & SHARP. M., 1973 - Morphology of Euclinostomum
heterostomum (Rudholphi, 1809) Trematode :
 Clinostomatidae from Bubuius ibis.
J. Helm., XLVII (1), 17-25.

DANDOTIA, M.R. & BHADAURIA, S. 1979 - On a new species of the
Brahamputrotrema Dayal et Gupta, 1954
 (Family-Lissorchiidae Poche 1926) from a
 fresh water fish, Puntius sophor. Riv.
Parassitol., 40 (1/2), 57-61.

DAS, B.K. AND RAHINULLAH, M. 1933 - Helminth parasites from
 certain fresh water fishes of India. Curr.
Sci., 1 (9), 278.

DAWES, B. 1946 - 'The trematoda', The Univ. Press. Cambridge;
 644 pp.

DAYAL, J., 1938 - On a new species of the genus Astiotrema
 Looss (1901) from the intestine of a fresh
 water fish Clarias batrachus from Lucknow.
Proc. Nat. Acad. Sci., 8(1), 10-14.

DAYAL, J., 1938 - Studies on the trematode parasites of fishes.
 A new trematode Neoganada barabankiae (non.
 gen., nov. spec.) from Clarias batrachus.
Ind. Acad. Sci. (Sec.B) 7, 132-137.

DAYAL, J., 1938 - Studies on the trematode parasites of fishes.

A new trematode Nizamia hyderbadi nov.gen.,
nov. sp. from the intestine of a fresh water
fish Ophiocephalus punctatus, Proc. Nat.
Acad. Sci., 8 (2), 53-58.

DAYAL, J., 1939 - On a new trematode Neoganada barabankiae
nov. gen. nov. sp. of the family Plagiorchiidae
from the intestine of a fish (Clarias batrachus).
(Abstract). Proc. 25th Ind. Sc. Congr., 6,
157-158.

DAYAL, J., 1949 - Trematode parasites of India fishes, Part II.
Indian J. Helminth., 1(2), 93-120.

DAYAL, J., 1950 - Trematode parasites of Indian fishes. Part III.
Two new trematodes of the family Allocreadiidae
from fresh water fishes of India. Ind. J.
Helm., 2(1), 1-10.

DAYAL, J. and Gupta, S.P., 1954 - A new trematode Brahamputro-
trema punctati n.g., n.sp. from a fresh water
fish Ophiocephalus punctatus (Bloch). Ibid, 190.

DAYAL, J. AND GUPTA, S.P. - 1964 - A description of four new
trematodes of the sub-family Haplorchiinae
(Looss, 1899) Poche, 1926 with a discussions
of systematics of the subfamily. Proc. 40th
Ind. Sci. Congr., 7, 188-191.

DEVRAJ, M. & RANGANATHAN, V. 1991 - Incidence of Isoparorchis hypselobagri (Billet, 1898) Trematoda, Hemiuroidae) among the cat fishes of Bhavani Sagar Reservoir. Indian J. Fish; 34, 1967: 232-250.

DUBININA AND BYCHOWSKY, 1954 - Trematodes of animals and man. (Skrjabin) vol. 9, 78-79.

DIETZ, E. 1909 - Die Echinostomiden der Vogel. Zool. Anz., 34 (6), 180-192.

DIETZ, E. 1910 - Die Echinostomiden der Vogel. Zool. Jahrb. supplm. 12 (3), 265-512.

DIRECTORATE OF FISHERIES, M.P., 1968 - A preliminary study of the Fish fauna of Madhya Pradesh.

DOLLFUS, R. Ph., 1932a - Trematodes de mammiferes, Oiseaux et poissons. Bull. Musc. natu. Hist., 4(5), 555-563.

DOLLFUS, R, Ph. 1932b - Trematodes result scient. Voyage Indes Orient. Neerl. Mem. Mus. Roy. Hist. Nat. Belg., hors ser 2(10), 13 pp.

DUBEY, G.P. & VERVA, M.N., 1968 - A preliminary study of the Fish Fauna of Madhya Pradesh.

DUDA, P.L. & VERMA, A.K., 1989 - Seasonal incidence of Gall bladder infection in the lacertilian hosts of Jammu.

DWIVEDI, M.P., 1965 - On a new species of the genus Genarchopsis Ozaki, 1925 from the stomach of Bufo melanostictus Schneid (Hemiuridae; Trematoda) Ind. J. Helm., XVII (1).

DWIVEDI, M.P., 1966 - On a new species of Brahmputrotrema Dayal et Gupta, 1954 from Phiocephalus punctatus. J. Zool. Soc. India. 21 (1), 133-136.

DWIVEDI, M.P. & CHAUHAN, B.S. 1970 - On some digenetic trematodes Part I. J. Zool. Soc. India. 21(1), (1969); 141 - 148.

DWIVEDI, M.P. & CHAUHAN, B.S. 1970 - On some digenetic trematodes Part II. J. Zool. Soc. India. 22, 87-96.

DWIVEDI, M.P. & CHAUHAN, B.S. 1970 - On some digenetic trematodes Part III. J. Zool. Soc. India 22 (1&2), 165-172.

EICHLER, W. 1949 - Some rules in ectoparasitism. Ann. Mag. nat. Hist., (12) 1, 588-598.

EJSMONT, L. 1932 - Note Sur le genre Isoparorchis. Ann. par., 10 (5), 453-457.

FISCHTHAL, J.H. & KUNTZ, R.E. 1963a - Trematode Parasites of fishes from Egypt. Part IV : A description of Monascus typicus (Odhner, 1911) (Fellodistomidae). Proc. helminth. Soc. Wash., 39(2), 177-182.

FISHTHAL, J.H. & KUNTZ, R.E. 1963b - Treematode parasites of fishes from Egypt. Part VI : The metacercaria of Euclinostomum heterostomum (Rudolphi 1809) Travassos, 1928 (Clinostomidae) with a review of the genus. Trans. Am. Microsc. Soc., 82(3): 335 - 342.

FISCHTHAL, J.H. & KUNTZ, R.E. 1963c - Trematode parasites of fishes from Egypt. Part VII : Orientocreadium betrachoides, Tubangui, 1931 (Plagiochioidea) from Clarias lazera, with a review of the genus and related form. J. Parasit., 49 (3) 451 - 464.

FISCHTHAL, J.H. & KUNTZ, R.E. 1963 - Trematode parasites of fishes from Egypt. Part V. Annotated record of some previously described forms. J. parasit., 49, 91 - 98.

FISCHTHAL, J.H. & KUNTZ, R.E. 1964 - Digenetic trematodes of fishes from Palwan Island, philippines, Part III families Hemiuridae and Lepocreadiidae. Ibid., 109 - 120.

FISCHTHAL, J.H. & THOMAS, J.D. 1968 - Digenetic trematodes of fresh water and marine fishes from Ghana, Proc. Helminth. Soc. Wash., 35(2) 126-140.

FISCHTHAL, J.H. & WILLIAMS, M.O. 1971 - Some digenetic trematodes of marine fishes from Sierra Leone. J. Helminth., 45(1), 45 - 50.

FISHER, R.A., 1944 - Statistical Methods for Research Workers. (9th Ed.) Oliver and Boyd, Edinoborg.

FOTEDAR, D.N., 1958 - On a new caryophyllaeid Cestode, *Adhenoscolex oreini* gen. et sp. Nov. from fresh water fish in Kashmir, and a note on some related genera J. Helminthol., 32, 1-16.

FOTEDAR, D.N., 1969 - Notes on some gorgoderid trematodes of fishes and amphibians in Kashmir and other parts of India and notes on the classification of the family Gorgoderidae Looss 1901, Kashmir Science. 6(1/2), 89-100.

GOHAR, N. 1934 - Les trematode parasites du milan egyptian, Milvus migrans avec description d' une espece et remarques sur less genres Haplorchis Looss et Monorchotrema Nishingori, 1924 Ann.Par., 12 (3), 213 - 227.

GUPTA, N.K. 1956 - Studies on the digenetic trematodes of marine fishes from the Gulf of Manner (India) Res. Bull. Punjab Univ. Sci., 89, 61-83.

GUPTA N.K. & MEHROTRA, V. 1970 - Observation on some already known digenetic trematodes of marine fishes of Ernakulam (South India). Res. Bull. Punjab Univ., Sci. 21, 87-99.

GUPTA, N.K. & SEHGAL, K.S. 1970 - Studies on some hemiurid trematodes from marine food fishes in India. Res. Bull. Punjab Univ. Sci., 21, 227-239.

GUPTA, N.K. & MUMARI, A. 1970 - On Hamacreadium manteri n.sp. (From : Allocreadiidae from fresh water fish Chela bacaila and chela gora at Ropar and Ludhiana (Abst.) J. Parasit., 56 (4, Sect.2) Int. Congr. Parasit (2nd) Washington, D.C., Sept. 6-12, 1970 Proc. Pt.I, 127-128.

GUPTA, N.K. & MUMARI, A. 1970 - On Chelatrema symthi n.g., n.Sp. (Trematoda: Hemiuridae Arnolineae) from a fresh water fish Chela bacaila from Ropar. (Abst.) Jour. Parasit, 56 (4, Sect.2) Int. Congr. Parasit. (2nd) Washington D.C. Sept. 6-12, 1970. Proc. Pt. I, 127.

GUPTA, N.K. & KUMARI, A. 1974 Publ. 1980 - On Hamacreadium Manteri n.sp. (Trematoda: Allocreadiidae) from a fresh water fish Chela bacaila and Chela gora at Ropar and Ludhiana Res. Bull.

Punjab. Univ. ^{Sci.} 25 (3 & 4), 199-202.

GUPTA, P.D. & SRIVASTAVA, C.B. 1960 - On Faustula Chauhani n.sp.
(Family-Fellodistomidae Nicoll, 1913). Ind.
J.Helm., 12 (2) 117-144.

GUPTA, S.P. 1950 - On a new trematode, Allocreadium thapari n.sp.
of the subfamily Allocreadiinae Looss, 1899,
from the intestine of fresh water fish, Rita
rita (Ham.). Ind. J. Helm., 2 (1), 17-22.

GUPTA, S.P. 1951 - Studies on the trematode parasites of food
fishes of U.P.A. new trematode cephalagonimus
heteropneustes n.sp. from a fresh water fish.
H. fossilis (Bloch). Indian J. Helminth., 3,
13-20.

GUPTA, S.P. 1951 - On a new trematode, Phyllodistomum singhai
n.sp. of family Gorgoderidae Looss 1899 from
intestine of fresh water fish. Mastacembalus
armatus (Ched.), Indian J. Helminth., 3, 21-28.

GUPTA S.P. 1951 - Three new trematodes of the family Hemiuridae
Luhe, 1901 from fresh water fishes of U.P.
Indian J. Helminth., 3(1), 41-54.

- GUPTA, S.P. 1951 - Studies on the trematode parasites of food fishes of U.P. Ind. J. Helm., 3(2), 101-108
- GUPTA, S.P. 1953 - Trematode Parasites of fresh water fishes, Ind. J. Helm., 5(1), 1-80.
- GUPTA, S.P. 1956 - Two new trematodes (Family - Allocreadiidae from the fresh water fishes of U.P. Ind. J. Helm., 8, 100-106.
- GUPTA, S.P. 1956 - Ind. J. Helminth., 8, 112 - 121.
- GUPTA, S.P. 1961 - A reference list of trematode parasites of fresh water fishes of India with a discussion on their systematic position Ind. J. Helm., 13, 35-60.
- GUPTA, S.P. 1961 - Caryophyllaeids (Cestoda) from fresh water fishes of India, Proc. Helminth. Soc. Wash., 28, 38-50.
- GUPTA, S.P. 1963 - On two new trematodes (Family-Allocreadiidae) Stossich, 1903) from the intestine of fresh water fishes of Banaras, U.P. Proc. Helm. Soc. Wash., 30, 96-100.

GUPTA, S.P. 1955 - Trematode Parasites of fresh water fishes,
Indian J. Helminth., 5(1) 1-80.

GUPTA, S.P. & AGARWAL V. 1968 - Pseudoparamacroderoides
Seenghali n.g., n.sp. (Allocreadiidae
wallininae) from the intestine of a fresh
water fish Mystus seenghala (Sykes) from
Lucknow, India. Indian J. Helminth.,
20 (1), 70-74.

GUPTA, S.P. & CHAKRAVARTHI, K.K. 1966 - On a treantode
Genarchopsis thapari n.sp. from the intestine
of a snake from Lucknow. Indian J. Helminth.,
18 (2), 177-180.

GUPTA, S.P. & ANIS JEHAN 1977 - A redescription of Euclinostomum
heterostomum (Rudolphi 1809) with a key to
the species of the genus Euclinostomum
Travassos, 1928. Ind.J.Helm. XXIX (2), 150-155.

GUPTA, V. AND PURI, M. 1979 Publ. 1980 - Four new species of
digenetic trematodes of fishes Ind. J. Helm.,
31 (1), 54 - 64.

GUPTA, V. AND PARNAR, S. 1984 - A on a new caryophyllaeid
Pseudocaryophyllaeus mackiewiczzi Sp.nov.,

from the intestine of a freshwater fish,
Heteropneustes fossilis (Bloch.) from
 Gorakhpur, U.P. Ind.J. Helminth., (1982, Pub.
 1984) 34, 136-138.

GUPTA, V. AND SINGH, S.R. 1984 - On a n.sp. Pseudocaryophyllaeus
ritai SP. Nov. (Family Caryophyllaeidae)
 from the intestine of a fresh water fish
Rita rita from river Gomati at Lucknow,
 U.P. Ind. J. Helminth., 35, 11-14.

GUPTA, V AND SINHA N. 1984 - On three new species of caryophy-
 llaeids from fresh water fishes of Lucknow.
Ind. J. Helminth., 36, 73-80.

HAFEZULLAN, M. & SIDDIQUI, A.H. 1970 - Digenetic trematodes of
 marine fishes of India. Part II, Fellodisto-
 midae. J. Parasit., 56 (5), 932-940.

HAFEEZULLAH M. 1971 - On some new and known digenetic trematodes
 from marine fishes of India. J. Helminth.,
 XLV (1), 73 - 88.

HAFIZUDDIN, A.K.M. & KHAN, H.R. 1973 - On a new species of the
 genus Genorchopsis Ozaki, 1925 (Digenea:
 Hemiuridae) Bangla-desh. J. Zool., 1, 107-110.

HANSON, M.L. 1950 - Some digenetic trematodes of marine fishes of Bermuda, Proc.HelminthSoc.Wash., 17(2), 74-88.

HANUMANTHA RAO, K. 1957- Ph.D. Thesis, Andhra University.

HINTER, G.W. III 1930 - Studies on the Caryophyllaeidae of North America, Illinois.Biol. Monograph, II (1927), 186 PP.

HOPKINS, S.H.E. 1949 - Some factors which have modified the phylogenetic relationships between the parasite and host in Mallophage. Proc.Linn. Soc. Lond., 161 (1), 37-38.

HUNTER, G.W. III- 1927 - Notes on the Caryophyllaeidae of North America.J. Parasitol., 14, 16-26.

JAIN, S P. & CHANDRA DEEP 1977 - Record of Some known metacercariae and adult trematodes of fishes from Agra district. Ind.J. Helm., XXIX (2), 84-86.

JAISWAL, G.P. 1957 - Studies on the trematode parasites of fishes and birds found in Hyderabad state I-IV. Zool. Jahrb. Syst., 85 (1-2), 1-12.

JAIWAL, G.P. 1967 - Investigation on the trematode fauna of the common food fishes of Hyderabad, A.P. Part I. A new species of Dorogenes Lühe, 1900 from a fresh water fish Channa (Ophiocephalus) Punctatus. Ind.J. Helm., Seminar Suppl. 18, 36-44.

JAIWAL, G.P. & NARAYAN, G. 1971 - Azygia Marulii Sp.n. (Trematoda) from a fresh water fish Channa marulius. Folia Parasit. 18 (2), 165-168.

James, R.HALL AND IVAN PRATT. 1969 - Some digenetic trematodes of Oregon Tidepool cottid fishes. J. Parasit. 55 (1), 201-205.

JOHNSTON, T.H. 1927 - New trematodes from an Australian siluroid. Trans & Proc. Roy. Soc. S. Austr. 51, 129-136.

JOHRI, G.N. 1959 - On a remarkable new Caryophyllaeid Oestode Hunterela mystei, mysteigen. et. sp. nov. from a fresh water fish in Delhi State. zeif. Parasitenkunde, 19, 368-374.

KAKAJI, V.L. 1968 - Studies on helminth parasites of Indian fishes. Part-I. Two trematode parasites of fresh water fishes from U.P. Indian. J. Helminth. XX(2) 136 - 144.

KAKAJI, V.L. 1969 - Studies on helminth parasites of Indian fishes. Part III. On Some species of the genus Allocreadium Looss (1900), Annals Parasit. Hum. Comp., 44 (2), 131-146.

KAKAJI, V.L. 1969 - Studies on helminth Parasites of Indian fishes. Part III. Some trematode parasites of fresh water fishes of U.P. Ind.J.Helm., XXI (1), 49 - 80.

KAW, B.L. 1944 - Studies on the helminth Parasites of Kashmir, Part III. Description on a new allocreadiid, Crepidostomum indicum from the Dal lake, Kashmir Proc. Indian Acad. Se., Sect. B, 20 (2), 72 - 77.

KAW, B.L. 1950 - Studies in Helminthology; Helminth Parasites of Kashmir. Part. I, Trematoda, Ind. J.Helm., 2 (2) 67 - 80.

KHALIL, L.F. 1961 - On a new trematode, Orientocreadium Lazeri SP nov. from a fresh water fish, Clarias Lazera in Sudan. J. Helm., 35, 259 - 262.

KHALIL, L.F. AND THURSTON, J.P. 1973 - Studies on the helminth Parasites of fresh water fishes of uganda, including the description of new species of digeneans. Revue de Zoologie et de Botanique Africaines, 87, 209 - 248.

- KAW, B.L. 1950 - Studies on the helminthology. Helminth
parasites of Kashmir Part I. Trematoda.
Indian J. Helminth., 2, 67 - 106.
- KUNDU, D.K. 1985 - On a new species of the genus Lytocestoides
Baylis, 1928 (Cestoidea : Caryophyllidea :
Lytocestidae) from a cobited fish,
Lepidocephalus guntea (Ham.), from West
Bengal. Bull. Zool. Surv. India, 7, 285-290.
- KUNDU, D.K., BHATTACHARYA, S.B. AND DATTA, I.B. 1985 -
Djombangia clariae n.sp. (Cestoidea :
Lytocestidae) from a siluroid fish in West
Bengal. Bull. Zool. Surv. India, 7, 151-154.
- LA RUE, G.R. 1957 - The classification of digenetic trematodes.
A review and a new system. Exp. parasit.,
6, 306 - 349.
- LEWIS, F.G. 1935 - The trematode genus Phyllodistomum Braun.
Trans. Amer. Micr. Soc. 54 (2), 103 - 117.
- LINTON, E. 1898 - Notes on trematode parasites of fishes
Proc. U.S. natn. Mus., 20 (1133), 507 - 548.

LINTON, E. 1934 - A new genus of trematodes belonging to the subfamily Allocreadiinae. J. Wash. Acad. Sci., 24 (2), 81 - 83.

LOOSS, A. 1907 - Zur kenntnis der Distomenfamilie Hemiuridae, Zool. Anz., 31, 585 - 620.

LUHE, M. 1901 - Uber Hemiuridae. Zool. Anz., 24, 394 - 488.

LYNSDALE, J.A. 1956 - On two new species of Lytocestus from Burma and the Sudan respectively. J. Helm., 30, 87 - 96.

MACKIEWICZ, J.S. 1963 - Subfamily status of Bovieniinae. Führmann, 1931 (Cestoidea : caryophyllaeidae) from catastomum commersoni (Lacepeda) (Pisces : Catastomidae) in North America. J. Parasitol. 51, 554-564.

MACKIEWICZ, J.S. 1965 - Isoglaridacris buylbocirrus gen. et sp.n. (Cestoidea : Caryophyllaeidae) from Catastomus commersoni in North America, J. Parasitol. 51, 554 - 564.

MACKIEWICZ, J.S. 1968 - Isoglaridacris hexacotyle Comb. n. (Cestoidea, Caryophyllidae) from catastomid fishes in Southwestern North America, Proc. helminth. Soc. Wash., 35, 193 - 196.

- MACKIEWICZ, J.S. 1972 - Caryophyllidea (Cestoidea) - a review.
Expl. Parasit.; 31, 417 - 512.
- MACKIEWICZ, J.S. 1981 - Synoptic review of the Caryophyllidae (Cestoidea) of India, Pakistan and Nepal.
Himalayan J. Sci., 1, 1 - 14.
- MACKIEWICZ, J.S. AND MURHAR, B.M. 1972 - Redescription of Bovienia serialis (Bovien, 1926) (Cestoidea: Caryophyllidea) from the cat fish Clarias batrachus (L.) in India. J. Helm., 46, 309-405.
- MADHAVI, R. 1978 - Life-history of Genarchopsis goppo, Ozaki, 1925 (Trematoda : Hemiuridae) from the fresh water fish Chann punctatus. J. Helm., 52, 251 - 259.
- MADHAVI, R. 1978 - Life history of Allocreadium fasciatusi Kakaji, 1969 (Trematoda : Allocreadiidae) from the fresh water fish Aplocheilus melastigma McCelland. J. Helm., 52, 51 - 59.
- MADHAVI, R. 1980 - Life history of Allocreadium handiai Pande, 1937 (Trematoda : Allocreadiidae) from the fresh water fish Channa punctatus (Bloch.). Z. Parasitenkd., 63, 89 - 97.

- MADHAVI, R. & H. RAO, K. 1977 - Anatomy of the female reproductive system in Allocreadium ophiocephali Srivastava, 1960 (Trematoda : Allocreadiidae) In Excerta. Parasitologion Memoria, 247-253.
- MANTER, H.W. 1940 - Digenetic trematodes of fishes from the Galapagos islands and the neighbouring Pacific. Allen Hancock Pacific exped., 2 (14), 329 - 454.
- MANTER, H.W. 1947 - The digenetic trematodes of marine fishes of Tortugas, Florida. Am. Midl. Nat., 38 (2), 257 - 416.
- MANTER, H.W. 1962 - Notes on the taxonomy of certain digenetic trematodes of South American fresh water fishes. Proc. Helm. Soc. Wash., 29(2), 97-102.
- MANTER, H.W. 1963 - Studies on the digenetic trematodes of fishes of Fiji. III Families Acanthocolpidae, Fellodistomidae and cryptogonimidae. J. Parasit., 49, 443 - 450.
- MANTER, H.W. 1965 - Host-parasite relationships. Proc. 26th Biol. Collicium 1965, Oregon State Univ. Publ., 59 - 71.

- MANTER, H.W. 1966 - A peculiar trematode, Gorgocephalus
Kyphosi gen. et sp. n. (Lepocreadiidae :
Gorgocephalinae subfam. n.), from a marine
fish of south Australia. J. Parasit. 52 (2)
347 - 350.
- MANTER, H.W. 1969 - Some digenetic trematodes of marine fishes
of New Caledonia. Part IV. Hemiuridae and
summary. Proc. Helm. Soc. Wash., 36 (2),
194 - 204.
- MANTER, H.W. & VAN CLEAVE, H.J. 1951 - Some digenetic trematodes
including 8 new species from marine fishes
of La Jolla, Calif. Proc. U.S. natn. Mus.,
101 (3279), 315 - 339.
- MANTER, H.W. & PRITCHARD, M.H. 1960a - Some hemiurid trematodes
from Hawaiian fishes. Proc. helminth. Soc.
Wash., 27 (1), 87 - 102.
- MANTER, H.W. & PRITCHARD, M.H. 1960b - Additional hemiurid
trematodes from Hawaiian fishes. Proc. helm.
Soc. Wash., 27 (2), 165 - 180.

MEHRA, H.R. 1930 - On a new species of caryophyllaeus O.F.

Muller from Kashmir with remarks on Lytocestus indicus (Moghe, 1925) Proc. 17th Sci. Cong., (3), 247.

MEHRA, H.R. 1957 - The classification of Digenetic Trematoda, Proc. Nat. Acad. Sci. Annual. number Bilock. Abstract : 71 - 72.

MEHRA, H.R. 1962 - Notes of the taxonomy of certain digenetic trematodes of South American fresh water fishes. Proc. Belm. Soc. Wash., 29(2), 97-102.

MEHRA, H.R. 1962 - Revision of Allocreadioidea Nicoll, 1934 Part I. Families : Lepocreadiidae Nicoll, 1934 Deropristidae n. fam., Homalometridae n. fam and Maseniidae Gupta, 1953. Proc. natn. Acad. Sci. India, 32(1), 1 - 22.

MEHRA, H.R. 1963 - Reivision of Fellodistomoidea (Syn.Fellodistomatoidea La Rue, 1957) : Families Fello-distomidae Woodcock, 1912, Gymnophallidae Dollfus, 1939, and Monodhelminthidae Dollfus 1937, Proc. Natn. Acad. Sci. India, 33(3B), 361 - 378.

- MEHRA, H.R. 1966 - Revision of Allocreadiidae Nicoll., 1934
Part II. Family Monorchidae Odhner, 1911
Asymphyloporidae Mehra, 1962 and Zoogonidae
Odhner, 1911, Publ. bv. author, 47 - 58.
- MESKAL, F.H. 1967 - Seasonal fluctuation in the population of
two common trematode species from the stomach
of the Cod. Sarsia, 26, 13 - 20.
- MOGHE, M.A. 1925 - Caryophyllaeus indicus n.sp. (Trematoda)
from the cat fish Clarias batrachus (Bl.)
Parasitology, 17, 232 - 235.
- MOGHE, M.A. 1931 - A supplementary description of Lytocetus
indicus Moghe. (Syn. caryophyllaeus indicus,
Moghe, 1925, Cestoda) Parasitology, 23, 84-87.
- MONNIG, H.O. 1926 - Helminthological notes. 11-12. Rept. Dir.
Vet. Educ. and Res., Dept. Agric. Union.
S. Africa (1), Sept., 221 - 228.
- MOTWANI, M.P. & SRIVASTAVA, C.E. 1961 - On the phyllodistomes
from the urinary bladder of siluroid fishes.
Ind. J. Helm., 13 (2), 93 - 99.

- MURHAR, B.M. 1963 - Crescentrovitus bioculus gen. nov., sp. nov., a fish cestode (Caryophyllaeidae) from Nagpur, India. Parasitology, 53, 413-418.
- NAGATY, H.F. 1942 - Trematodes of fishes from the Red Sea. III. On seven new Allocreadiid Species. Public. Mar. Bio. Stat. Ghardaqui (Red Sea) 4, 27 :
- NICOLL, W. 1914a - Trematode parasites from animals dying in the Zoological Society's Gardens during 1911-1912. Proc. Zool. Soc. Lond., 1, 139-154.
- NICOLL, W. 1914b - The trematode Parasites of North Queensland. I. Parasitology, 6, 333 - 350.
- NICOLL, W. 1915 - The trematode parasites of North Queensland III - Parasites of fishes, Parasitology, 8 (1), 22 - 41.
- NISHIGORI, M. 1924 - Two new trematodes of the family Heterophyidae found in Formosa. J. Med. Assoc. Formosa, 237.
- NIYOGI, A., GUPTA, A.K. AND AGARWAL, S.M. 1982 - Morphology of Lucknowia indica sp. nov. (Lytocestidae : Caryophyllidea), Proc. Indian. Acad. Parasit., 3, 17 - 22.

NYBELIN, O. 1926 - Zur Helminthan fauna der Susswasserfische.

Schwedens, I. Phyllodistomen. Got., Kungl.

Vitt. Samh. Handl. 31 (3), 29.

OZAKI, Y. 1925a - Preliminary notes on a trematode with annus.

J. Parasit. 12, 51 - 53.

OZAKI, Y. 1925b - On a new genus of fish trematode, Genarchopsis

Jap. J. Zool., 1(3), 101 - 108.

PANDE, B.P.

The trematode genus Allocreadium in Indian fresh water fishes. Proc. Indian Acad Sci., 7, 54 - 60.

PANDE, B.P. 1934 - On a new trematode from the Indian fresh

water fish. Proc. Nat. Acad. Sci. India, 4(1), 107 - 112.

PANDE, B.P. 1937 - Two new fish trematodes from Allahabad.

Proc. Nat. Acad. Sci., 7(2) 111 - 115.

PANDE, B.P. 1937 - Morphology and relationships of a new

digenetic trematode from an Indian fresh water fish. Ann. Mag. Nat. Hist. Ser., 10 (20), 415 - 421.

PANDE, B.P. 1938 - The trematode genus Allocreadium in North

Indian fresh water fishes. Proc. Ind. Acad. Sci., 7 (2), 54 - 60.

- PANDE, B.P. 1938 - On two new trematodes from Indian Cyprinoid fishes with remarks on the genus Allocreadium Looss, Proc. Nat. Acad. Sci. India, 8 (4), 110 - 115.
- PARUKHIM A.M. AND SYSA, V.N. 1975 - The incidence of infection in Notothenioid fish in Sub-Antarctic water. Problmy Parazitologii. 'Navkova Dumka' 97 - 99.
- PARK, J.T. 1939 - Fish trematodes from Tyoson II. Some new digenetic trematodes from marine fishes. J. Med., 10 (1) 7 - 18.
- PATHAK, A. & THAKUR, J. 1989 - Studies on Helminth infestation in human Population of Bhopal (India.)
- PEARSON, J.C. 1964 - A revision of the subfamily Haplorchinae Looss, 1899 (Heterophyidae) I. The Haplorchis group. Parasit, 54 (4), 601 - 676.
- PEREZ VIGUERAS, I. 1955 - Contribucional conocimientos de la fauna helminthologica cubana. Mem. Soc. Cub. Hist. Nat., 22 (1), 21 - 71.
- PERSHAD, R.S. 1965 - On a new species of the genus Rhynchocreadium Srivastava (1962) from the fresh water fish, Rhynchobdella aculeata in Hyderabad, A.P. Zool. Anz., 175, 4-6.

- PETRUSHEVSKI, G.K. & PETRUSHEVSKAJA, M.G. 1960 - The accuracy of quantitative indices under the study of parasito-fauna of fishes - paras. Sobr. Zool., 19, 333 - 343.
- FIGULEWSKY, S.W. 1931 - Neue Arten Trematoden aus Fischen des Dnije prbassins. Zool. Anz., 104, 8-18.
- PANDE, B.P. AND SHUKLA, R.P. 1969 - Studies on the metacercaria of fresh water fishes of India VI. On the morphology of metacercaria of Isoparorchis hypselobagri (Billet, 1898) with a note on its development Proc. Nat. Acad. Sci. India, 39, 185 - 190.
- PANDE, B.P. AND SHUKLA, R.P. 1976 - Haplorchoides Chen, 1949 (Haplorchinae : Heterophyidae) in fresh water fishes. J. Helm., 50 (3) 181 - 192.
- PANDEY, K.C. 1971 - Studies on trematode parasites of fishes of Lucknow (India) I. Proc. Nat. Acad. Sci. India, 41 (B) III, 311 - 314.
- PANDEY, K.C. 1973 - Studies on some known and unknown trematode Parasites. Ind. J. Zootomy, 14 (3), 197 - 219.

- PANDEY, K.C. 1973 - A restudy of Genarchopsis goppo (Tubangui) Ozaki, 1925 with a note on validity of certain species. Ind.J. Zootomy, 14 (3), 167 - 174.
- PANDEY, K.C. 1973 - Studies on some cestodes from fishes, birds and mammals. Ind. J. Zool., 15, 221 - 226.
- PANDEY, V. 1979 - Metacercaria and adult of Haplorchoides attenuatus infesting local fishes. Ind. J. Anim. Sci., 49 (4), 303 - 307.
- PRICE, E.W. 1934 - New digenetic trematodes from marine fishes. Smithson. Misc Coll., 91, 8 pp.
- PURDHOO, S. 1944 - On some trematodes from ceylon. Ann. Mag. Nat. Hist., 11-S (73), 11, 1-13.
- RAHIMULLAH, M. AND DAS, B.K. 1934 - On the trematode infection in certain Indian fishes. Curr. Sci., 3 (5), 206 - 207.
- RAI, P. 1970 - A redescription of Allocreadium mehrai Gupta, 1956 and A. handiai Pande, 1937 with a key to identification of species of Allocreadium

Looss, 1900 from fresh water fishes of India. Ind. J. Sci. Indust. (B) (4(1) 35 - 40.

RAI, P. 1971 - On the genus Genarchopsis Ozaki (Trematoda : Hemurida) from fresh water fishes of Indian. Agra Univ. J. Res., 20, 27 - 34.

RAI, P AND PANDE, B.P. 1965 - Isoparorchid infection in some hitherto unrecorded fish species. Curr. Sci., 34, 586 - 588.

RAI, P AND PANDE, B.P. 1967 - Observation on the haplorchine flukes hitherto known from Indian siluroid fishes, Ag. Univ. J. Res. Sci., 16, 17 -24.

RAI, P. AND PANDE, B.P. 1975 - Isoparorchid infection in some hitherto unrecorded fish species. Curr. Sci., 34, 586,- 588.

RAI, S.L. 1961 - Observations on the life history of Phyllodistomum Srivastava n.sp. (Trematode : Gorgorediae). I. Metarcercaria and experimental adult. Proc. Nat. Acad. Sci. India, annual No. 111 - 112.

- RAI, S.L. 1962 - Studies on three new species of the genus Allocreadium Looss, 1900 from the intestine of Barbus tor (Ham.), Parasit., 52, 23 - 30.
- RAI, S.L. 1964 - Observations on the life history of P. Srivastavi sp.n. (Gorgoderidae). Parasit., 54, 43 - 51.
- RAZVI, S.S.H. 1971 - Digenea of Pakistan fresh water fishes. I. Haplorchoides indicus n.sp. (Family - cryptogonimidae) from the intestine of Mystus aor (Ham.) Sukkur, West Pakistan, Pak. J. Sci. Indu. Res., 14, 518 - 520.
- SAHAY, S.N. AND SAHAY, U. 1977 - On a new caryophyllid cestode, Djombangia caballeri sp. nov. from fresh water fish Heteropneustes fossilis in chotanagpur with an amendment of the generic character. Excerpta parasitologica en Memoria del Doctor Eduardo Caballero Y Caballerao, Mexico, 4, 371 - 376.
- SAKSENA, J.N. 1958 - On a new species of the genus Allocreadium Looss, 1900, of the family Allocreadiidae stossich, 1903 from the intestine of Mastacembelus armatus. Proc. Nat. Acad. Sci., 28 (3), 227 - 230.

SAKSENA, J.N. 1958 - Studies on two new species of the genus Orientocreadium (Trematoda : Allocreadiidae) from the intestine of Clarias magur Proc. Nat. Acad. Sci., (B) 28 (1), 58 - 64.

SAKSENA, J.N. 1960 - Studies on a new species of the genus Orientocreadium (Trematoda : Allocreadiidae) from the intestine of Clarias Magur. Proc. Nat. Acad. Sci., (B), 30 (1), 83 - 86.

SAKSENA, J.N. 1965 - An analysis of the trematode genus orientocreadium Tubangui, 1931 . Proc. Nat. Acad. Sci., (B) XXXV (IV), 367 - 370.

SATPUTE, L.R. AND AGARWAL, S.M. 1974 - Diverticulosis of the fish duodenum infested with cestodes. Indian J. Exp. Biol., 12, 373 - 375.

SATPUTE, L.R. & AGARWAL, S.M. 1980 - Introvertus raipurensis gen. Nov., a fish cestode (Cestoda : Caryophyllidea: Lytocestidae) from Raipur, India, Proc. Ind. Acad. Parasit. 1, 17 - 19.

SCOTT, J.S. 1975 - Incidence of trematode parasites of American Plaice Hippoglossoides platessoides of the scotian shelf and Gulf of St. Lawrence in relation to fish length and food. J. Fisheries Research Board Canada, 32 (4), 479 - 482.

- SIDDIGI, A.H. & NIZAMI, W.A. 1988 - Incidence of Isoparorchis hypselobagri Billet, 1898 (Trematoda : Isoparorchidae) in Wallago attu with remarks on its life cycle Acta. Parasitologica Polonica, 35 (21/35), 223 - 227.
- SIMEY, P.H. 1929 - Fish trematodes of the lower Talahatchie River. Am. Midl. Nat., 11 (12), 563 - 602.
- SIMHA, S.S. & PERSAD, R.S. 1964 - Description and observation on the development of Azvgia asiatics n.sp. from fresh water fish O.punctatus. Rev. Parasit., 25, 25 - 30.
- SINGH, K.S. 1953 - Echinostoma thapari n.sp., from an Indian fish, Notopterus chitala (Ham.) Thapar Comm. Vol., 245 - 250.
- SINGH, K.S. 1954 - Some trematode collected in India. Trans. Amer. Micros. Soc., 73, 202 - 210.
- SINGH, K.S. 1957 - Diplostomilum elongatus n.sp. (Trematoda) from a fresh water fish Trichogastar fasciatus from India. J. Parasit., 43, 371 - 373.
- SINGH, S.P. AND SINHA, D.P. 1975 - Two new trematodes from fresh water fishes of Bihar. Dr. B.S. Chauhan Comm. Vol., 411 - 417.

SINGH, S.P. AND SINHA, D.P. 1976 - On Rhipidoeotyle Vachius sp. n. (Bucephalidae : Gasterostomata : Trematoda) from an Indian fresh water fish Eutropiichthys vacha. Annals Zool., XII (4).

SINGH, S.P. AND SINHA, D.P. 1977 - On a new gasterostome trematode Bucephalus tetratentacularis n.sp. from an Indian fresh water fish. Sciaena Coiter, Ceylon J. Sci., 12(2), 115 - 118.

SINGH, S.P. AND SINHA, D.P. 1980 - The occurrence of some trematodes in fresh water fishes of Bihar, new host records. Ind. J. Anim. Res., 14 (1) 57 - 62.

SINHA, A., SAHAY, V. & PRASAD, D. 1974 - Studies on the Digenetic trematodes of Bihar - II Part. Ind. Jour. Animal. Research Haryana (India), 8 (1), 39 - 41.

SINHA, S.S. & RASHEED, U. 1981 - An anomaly in caryophyllid cestode, Lytocestus indicus moghe, 1931. Ind. J. Parasitol., 197 - 198.

SKRJABIN, K.I. 1947 - Trematodes of Animals and Man. Moskov, 1, 515 pp.

SKRJABIN, K.I. 1954 - Trematodes of Animals and Man.

Moskov, 9, 654 pp.

SKRJABIN, K.I. 1961 - Trematodes of Animals and Man.

Moskov, 19, 1 - 192.

SKRJABIN, K.I. 1962 - Trematodes of Animal and Man.

Moskov, 20, 563 pp.

SKRJABIN K.I. & ANTIPIN, D.N. 1956 - In K.I. SKrjabin

Trematodes of Animal and Man. Moskov, 12,
53 - 932.

SKRJABIN, K.I. & GUSHANSKAYA, L.KH. 1955 - In K.I. Skrjabin

Trematodes of Animals and Man. Moskov,
10, 622 - 634.

SKRJABIN, K.I. & KOVAL, B.P. 1956 - In K.I. Skrjabin Trematodes

of Animals and Man Moskov, 13, 165 - 452.

SKRJABIN, K.I. & PETROV, V.P. 1958 - In K.I. Skrjabin :

Trematodes of Animals and Man. Moskov, 15
320 - 424.

SOGANDARES BERNAL, F & HUTTON, R.F. 1959 - Studies on the

helminth parasites from the coast of Florida

III. Digenetic trematodes of marine fishes

from Tampa and Boca ciegu Bays. Hain, 45,

337 - 446.

SOOTA, T.D. & GHOSH, R.K. 1977 - On some trematodes from
Darjiling, West Bengal. Rec. Zool. Surv.
India, 73, 223 - 228.

Southwell, T. 1913 - Notes from the Bengal Fisheries Laboratory
Indian Museum, No.1 Rec. Ind. Mus., 9, 79-103.

Southwell, T. AND PRASAD, B. 1918 - Notes from Bengal Fisheries
Laboratory, No. 5, Parasites of Indian
fishes, with a note on car cinoma in the
climbing perch. Rec. Indian Mus., 15 (5),
341 - 355.

SPARKS, A.R. 1958 - Some digenetic trematodes of fishes of
Grand Isle. Louisians. Proc. La. Acad. Sci.,
20, 71 - 82.

SRIVASTAVA, C.B. 1962 - On Pycnadena komiyai n.sp. (Trematoda :
Allocreadiidae, J. Med. Sci. Bio., 15,
275 - 277.

SRIVASTAVA, C.B. 1962 - A new Allocreadiid trematode,
Rhynchocreadium aculeata gen. et. sp. nov.
from the fresh water eel, Rhynchobdella
aculeata (Bloch), Ind. J. Helm., 14, 1 - 4.

- SRIVASTAVA, C.B. 1963 - On Falliorchis vermai sp. nov.,
(Trematoda : Allocreadiidae) from
Eutropiichthys vacha (Ham.). Ind. J. Helm.,
15, 45 - 47.
- SRIVASTAVA, C.B. 1972 - Studies on the Digenetic Trematodes
of vertebrates. Ph.D. Thesis. Calcutta Univ.
- SRIVASTAVA, C.B. 1980 - Estimation of Helminthic infections
Proc. workshop Tech. Parasitol. Zool. Surv.
India, 29 - 31.
- SRIVASTAVA, C.B. 1981 - Some observation on the Population
fluctuation of Prosorhynchoides magnum
(Verma, 1936) in the fish Silonia silonia
(Ham.) Proc. Symp. Ecol. Anim. Popul. Zool.
Surv. India, Pt. 2, 133 - 136.
- SRIVASTAVA, CB. & CHAUHAN, B.S. 1972 - A review of Indian
Gasterostomes (Trematoda) Rec. Zool. Surv.
India, 67, 1 - 13.
- SRIVASTAVA, C.B. & GHOSH, R.K. 1967 - On a new genus
Paramacrolecithus (Trematoda : Allocreadiidae)
from a Cyprinifish, Rasbora rasbora (Ham.)
Proc. Ind. Acad. Sci., 66, 179 - 183.

- SRIVASTAVA, C.B. & MUKERJEE, G.D. 1988 - Studies on the incidence of infestation of Isoparorchis hypselobagri (Billet, 1898) metacercaria in two species of fishes of the genus Mystus. J. Zool. Soc. India, 38 (1/2), 131 - 137.
- SRIVASTAVA, C.B. & SINGH, S.P. 1967 - On Eucreadium jhingrani n.sp. (Trematoda : Allocreadiidae) Proceedings of the Nat. Acad. of Sci., India, section B, Vol. XXXVII, Part I, 1967 pp. 117 - 119.
- SRIVASTAVA, H.D. 1933 - On new trematodes of frogs and fishes of the united provinces, India. Part I, Distomes of the family Hemiuridae from North Indian fishes and frogs with a systematic discussion on the family Halipegidae and the genera Vitellotrema Guberlet and Genarchopsis Ozaki. Bull. Acad. Sci. Allahabad., 3 (1), 41 - 60.
- SRIVASTAVA, H.D. 1934 - On new trematodes of frogs and fishes of the united provinces, India. Part IV. The occurrence and seasonal incidence at infection of certain trematodes in the above hosts. Bull. Acad. Sci. U.P. Agra and Oudh., 4 (1) 113 - 119.

SRIVASTAVA, H.D. 1935 - New hemiurids (Trematoda) from Indian fresh water fishes. Part I. New distomes of the genus Lecithaster Luhe, 1901, from Clupea ilisha. Proc. Acad. Sci. India, 4 (4) 381 - 387.

SRIVASTAVA, H.D. 1935 - New parasites of the genus Orientophorus n.gen. (Family Fellodistomidae) from an Indian fresh water fish, Clupea ilisha. Parasit; 27 (3), 374 - 382.

SRIVASTAVA, H.D. 1935 - New hemiurids (Trematoda) from Indian fresh water fishes. Part II. A rare parasite of the subfamily Dinurinae from Clupea ilisha. Zeitschr. Par, 8(1), 135 - 138.

SRIVASTAVA, H.D. 1935 - Studies on the family Heterophyidae Odhner, 1914 Part II. Four new parasites of the genus Haplorchis Looss, 1899, from Indian fresh water fishes with a revision of the genus. Proc. Acad. Sci. India, 5 (1), 75 - 85.

SRIVASTAVA, H.D. 1935 - New hemiuridae from Indian fresh water fishes II. A rare parasite of the sub family Binurinae from Clupea ilisha. Z. Par, 8, 135-138.

- SRIVASTAVA, H.D. 1936 - A rare parasite of the family Monorchudae Odhner, 1911, from an Indian fresh water fish, Ophiocephalus. Ann. Mag. Nat. Hist., 10, S (98) V. 17, 319 - 324.
- SRIVASTAVA, H.D. 1937 - New hemiurids (Trematoda) from Indian marine food fishes. Part III : Two new parasites of the genus Lecithocladium Looss from Indian food fishes with a revision of the genus. Proc. 24th Indian Sci. Congr., 295 - 296.
- SRIVASTAVA, H.D. - 1937 - New Allocreadiids (Trematoda) from Indian marine food fishes. Part III. Pendunculacetabulum Pedicellata n.sp. from the gut of Chiloscyllium indicum. Proc. 24th Ind. Sci. Cong., 298.
- SRIVASTAVA, H.D. 1938 - Studies on the amphistomatous parasites of Indian food fishes. Part I. Two new genera of amphistomes from an Indian fresh water fish, silundia gangatica (Cuv. d. Val.). Ind. J. Sci. Anim. Husb., 8 (4), 367 - 374.

SRIVASTAVA, H.D. 1938 - A new gorgoderid trematode from the urinary bladder of an Indian migratory fish, Belone Strongylurg. Ind. J. Vet. Sci. Ani. Husb., 8(4), 391 - 393.

SRIVASTAVA, H.D. 1938 - Studies on Gasterostomatous parasites of Indian food fishes (with ten text figs.). Ind. J. Vet. Sci., 8, 317 - 340.

SRIVASTAVA, H.D. 1939 - Studies on the family Heterophyidae Odhner, 1914, Part III, Parasites belonging to a new subfamily polyrchitreminae from an Indian fresh water fish. Ind. J. Vet. Sci. Anim. Husb., 9 (2), 165 - 168.

SRIVASTAVA, H.D. 1941a - New Fellodistomids (Trematoda) from Indian fishes. Part III a new parasite of the genus Haplocladus Odhner, 1911 from an Indian marine fish. Indian J. Vet. Sci., II, (1), 42 - 44.

SRIVASTAVA, H.D. 1941b - New Allocreadiids (Trematoda) from Indian marine food fishes Part V : A new parasite of the genus Lepocreadioides Yamaguti 1936. Ind. J. Vet. Sci., 11 (1), 52 - 54.

- SRIVASTAVA, N.N. 1951 - A new digenetic trematode, Eumassenia moradabadensis n.g., n.sp. (Plagiorchiidae Luhe, 1901 : Sub family Maseniinae (Chhaterji, 1933) from a fresh water fish, Heteropneustes fossilis, Ind. J. Helm., 3 (1) 1 - 6.
- SRIVASTAVA, O.N. 1950 - A preliminary note on the life cycle of Clinostomum sp. and Euclinostomum heterostomum (Rudholphi, 1809). Proc. Nat. Acad. Sci. India, (Sect. B) 20, 85 - 92.
- SRIVASTAVA, P.S. 1960 - On a new trematode Allocreadium ophiocephali n.sp. of the family Allocreadiidae stossich, 1903 from the Intestine of Ophiocephalus punctatus. Ind. J. Helm., 12(2) 108 - 113.
- STAFFORD, J. 1904 - Trematodes from Canadian fishes. Zool. Anz. 37 (16-17), 481 - 495, Also in Zool. ctbl. 13 (1-2), 42 - 43 (1906).
- STEEN, E.B. 1938 - Two new species of Phyllodistomum (Trematoda-Gorgoderidae) from Indians fishes. Am. Midl. Nat., 20 (1), 206 - 210.

STEEN, E.B. 1938 - A note on a phyllodistomid trematode from the urinary bladder of the Miller's thumb, cottus sp. Proc. Ind. Acad. Sci. 47, 228,

STUNKARD, H.W. 1930 - Another trematode with two anal openings Anat. Rec., 47 (3), 363 p.

STUNKARD, H.W. 1931 - Further observations on the occurrence of anal openings in digenetic trematodes. Parasitkde., 3, 713 - 725.

THAPAR, G.S. 1936 - Parasitic worms and disease. Lucknow Univ. Stud., No. III, : 46 pp.

THAPAR, G.S. 1956 - Helminthology. In Progress of Science Sec. 7 : Zoology. Nat. Inst. Sci. India Publ., 1 - 18.

THAPAR, G.S. & DAYAL, J. 1959 - Proc. Nat. Acad. Sci. India, 29, 34 - 36.

TRAVASSOS, L. 1928 - Sur La systematique de la famille des Clinostomidae Luhe, 1901, Compt. Rend. Soc. Biol. Paris., 98, 643 - 644.

TUBANGUI, M.A. 1931 - Trematode Parasites of Philippine vertebrates III. Flukes from fish and reptiles. Philipp. J. Sci., 44 (4) 417 - 423.

- TUBANGUI, M.A. 1933 - Trematode Parasites of Philippine vertebrates. VI, Descriptions of new species and classification. Philipp. J. Sci., 52 (2) 167 - 197.
- TUBANGUI, M.A. & MASILUNGAN, V. 1935 - Trematode parasites of vertebrates VII. Additional records of new species. Philipp. J. Sci., 58 (4), 435 - 445.
- VAN CLEAVE, H.J. & MUELLER, J.F. 1932 - Parasites of the oneida lake fishes, Part I, Descriptions of new species and some general considerations, especially concerning the trematode family Heterophyidae. Roosevelt wild Life Ann., 3 (2), 79 - 137.
- VASANTHA KUMARI, M. 1972 - A new species of the genus Monascus Looss, 1907 (Trematoda : Digenea : Fellodistomidae) Parasitic on flat fishes. J. Zool. Soc. India, (In press)
- VASANTHA KUMARI, M. 1973 - Studies on Helminth parasites Ph.D. Thesis Calcutta University, Calcutta.
- VANDER KUYP, E. 1953 - Euclinostomum heterostomum infcelebes. Doc. Med. Geogr. et Trop., 5, 329 - 332.
- VENKATANARSAIAH, J. 1979 - Gyrodactylus hyderabadensis n.sp. from Channa sp. from Hyderabad. Ind. J. Par., 3(1), 41-42.

- VELASQUEZ, C.C. 1962 - Some hemiurid trematodes from
Philippine fishes. Parasit., 48 (4), 539-544.
- VERMA, S.L. 1971 - Helminth parasites of fresh water fishes,
Part I. on two caryophyllaeids from fresh
water fishes of Lucknow. Indian J. Helminth.,
23, 71 - 80.
- VERMA S.L. 1973 - Helminth parasites of fresh water fishes.
Part II. On some trematode parasites from
fresh water fishes of Lucknow, Proc. Nat.
Acad. Sci. India, (Sec. B) 43 (1/2), 81 -, 86.
- VERMA S.L. 1974 - Helminth parasites of fresh water fishes,
Part X. On two new trematode parasites of
fresh water fishes of Lucknow (India). Rev.
Biol. Trop., 21, 181 - 186.
- WITENBERG, G.G. 1929 - Studies on the trematode family
Heterophyidae. Ann. Trop. Med. Par., 23 (2)
131 - 239.
- WITENBERG, G.G. 1930 - Corrections to my paper 'Studies on
the trematode family Heterophyidae'. Ann.
Mag. Nat. Hist., 10 S 28, 5, 412 - 414.

- WOODLAND, W.N.F. 1923 - On some remarkable new forms of Caryophyllaeidae from the Angloegyptian Sudan, and a revision of the cestodaria. Jr. Microsc. Sci., (N.S.) 67, 435 - 472.
- WOOLCOCK, V. 1935 - Digenetic trematodes from Australian fishes (With plates VIII and IX and 6 figures in the text). Parasitology, 27 (3), 309 - 331.
- MU, K. 1938 - On the occurrence of Isoparorchis amongst fishes in China (Trematoda : Isoparorchidae). Peking Nat. Hist. Bull., 12 (4), 273 - 277.
- YAMAGUTI, S. 1934 - Studies on the helminth fauna of Japan, Part 2, Trematodes of Fishes. 1. Jap. J. Zool. 5 (3), 249 - 541.
- YAMAGUTI, S. 1938 - Studies on the Helminth fauna of Japan. Part 21, Trematodes of fishes IV. Publ. by the author, 139 pp.
- YAMAGUTI, S. 1939 - Studies on helminth fauna of Japan, Part, 26, Trematodes of fishes VI. Jap. J. Zool., 8 (2), 211 - 230.
- YAMAGUTI, S. 1940 - Studies on the helminth fauna of Japan, Part, 31, Trematodes of fishes VII. Jap. J. Zool., 9, 36 - 108.

YAMAGUTI, S. 1953 - Parasitic worms mainly from celebes,
Part III Digenetic trematodes of fishes II.
Acta. Med. Okavama, 8(3), 257 - 295.

YAMAGUTI, S. 1954 - 'Systema Helminthum' Part I. Digenetic
trematodes of fishes. Published be author.
Tokyo (1953). 405 p.

YAMAGUTI, S. 1958 - 'Systema Helminthum'. Vol. I. The
digenetic trematodes of vertebrates, Part
I & II, Interscience Publishers, New York.

YAMAGUTI, S. 1971 - 'Synopsis of digenetic trematodes of
vertebrates'. Vol. I & II. Keigaku Publishing
Co., Japan.

- - - - -